CHAPTER 4.5.

IMPROVING POST-FACTORY PERFORMANCE OF STERILE MALE FRUIT FLIES IN SUPPORT OF THE STERILE INSECT TECHNIQUE

R. PEREIRA¹, B. YUVAL², P. LIEDO³, P. E. A. TEAL⁴,
T. E. SHELLY⁵, D. O. MCINNIS⁴, I. HAQ⁶, P. W. TAYLOR⁷
AND J. HENDRICHS¹

¹*Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture,
International Atomic Energy Agency, A-1400 Vienna, Austria*

Email: R.Cardoso-Pereira@iaea.org

²*Department of Entomology, Hebrew University of Jerusalem, Israel*

³*El Colegio de la Frontera Sur (ECOSUR), Tapachula, Chiapas, Mexico*

⁴*Deceased*

⁵*USDA/APHIS, Waimanalo, HI, USA*

⁶*National Agricultural Research Centre, Park Road, Islamabad, Pakistan*

⁷*Applied Biosciences, Macquarie University, Sydney, Australia*

TABLE OF CONTENTS

1. INTRODUCTION	632
2. NUTRITIONAL SUPPLEMENTS.....	633
3. HORMONAL TREATMENTS	636
4. SEMIOCHEMICAL TREATMENTS	639
5. CONCLUSIONS	643
6. REFERENCES.....	644

Pages 631–656

V. A. Dyck, J. Hendrichs and A. S. Robinson (eds.), Sterile Insect Technique. Principles and Practice in Area-Wide Integrated Pest Management. Second Edition.

Published with the permission of © 2021, US Government. CRC Press, Boca Raton, Florida, USA.

SUMMARY

The sterile insect technique (SIT) is being applied against tephritid fruit fly pests in many areas of the world. Currently, fruit fly factories have the capacity to produce and sterilize several billion (thousand million) sterile insects per week, and to make them available for shipment to their final destinations. At sterile fly emergence and release facilities, the emerging flies are fed and held close to maturity, and then collected for area-wide release. While much research effort has been invested in improving mass-rearing and quality-control procedures at the fly-factory level, the post-factory handling of sterile flies has received much less attention. However, research (conducted mainly from 2000 onwards) has focussed on developing and validating ways of improving sterile male performance through better management during a critical period (starting with the arrival of pupae at the fly emergence and release facility and ending with the release of the sterile flies in the field). This chapter summarizes the progress made on this subject for fruit fly species (in the genera *Anastrepha*, *Bactrocera*, *Ceratitis* and *Zeugodacus*) against which the SIT is being applied. To increase the effectiveness of fruit fly SIT programmes, exposure of sterile males to nutritional, hormonal, and semiochemical treatments has been assessed for improvement in sterile male performance, and enhancement of post-factory handling and release methods. Incorporation of protein and juvenile hormone into pre-release *Anastrepha* spp. diets significantly accelerates sterile male maturation and improves sexual performance in several species. Improved or probiotic adult diets and semiochemical treatments using ginger root oil or citrus oils in *Ceratitis capitata*, and methyl eugenol and raspberry ketone in *Bactrocera* and *Zeugodacus* species, significantly increase sterile male mating competitiveness. Some of these treatments and improvements have been transferred to, and are being applied routinely in, operational programmes. However, these efforts need to be further strengthened to assess the interaction among different environmental and holding conditions, treatments and release systems, and to improve further the performance of mass-produced sterile males, a critical component of increasing the effectiveness of operational programmes.

1. INTRODUCTION

Application of the sterile insect technique (SIT), as part of an area-wide integrated pest management (AW-IPM) approach against fruit fly (Diptera: Tephritidae) pests, is gaining momentum, with active programmes targeting fruit fly pest species of economic importance in North, Central and South America, Europe, Middle East, Asia, Africa, and Australia (Enkerlin, this volume). There are now fruit fly factories with the capacity to mass-rear several billion (thousand million) sterile male insects per week (for detailed information, note the World-Wide Directory of SIT Facilities (DIR-SIT 2020)) (Parker, Mamai et al., this volume). After irradiation (Bakri et al., this volume), pupae are shipped to their destination, where they are processed in fly emergence and release facilities (FAO/IAEA 2017; Dowell et al., this volume). Several days after emergence, feeding, and sexual maturation, the sterile flies are released in the field, where the sterile males are expected to mate with, and transfer an effective ejaculate to, wild females. An industrial process, consisting of numerous complex steps, is required to achieve a biological goal (Pereira et al. 2013a).

Several quality-control and quality-assurance protocols have been developed to ensure that the released flies fulfil a series of minimum quality standards (FAO/IAEA/USDA 2019; Parker, Vreysen et al., this volume), but exigencies of the industrial process often affect the biological qualities of the final product.

The mass-rearing of fruit flies up to the pupal stage, including sterilization, takes place in specially designed insect factories (where insects are adapted to indoor conditions) (FAO/IAEA 2018). As late-stage pupae, the mass-reared insects are irradiated and shipped to fly emergence and release facilities, where they are placed in containers/release bags, boxes or towers for adult emergence, feeding and holding

before they are released into target field areas (FAO/IAEA 2017; Dowell et al., this volume). During this critical period at these facilities, there is the potential to manipulate sterile males in a manner that will improve their mating success in the field (Pereira et al. 2013a). This includes optimizing the provision of nutritional supplements (Yuval et al. 2002), and hormonal (Teal et al. 2000) and semiochemical treatments (Shelly 2001), as well as the holding conditions and preparations for release (Tween and Rendón 2007; USDA/APHIS 2009; FAO/IAEA 2017).

An FAO/IAEA Coordinated Research Project (CRP) on “Improving Sterile Male Performance in Fruit Fly SIT Programmes” was conducted (2004 to 2009) with the objective of increasing the effectiveness of the SIT by improving the performance of sterile males after mass-rearing. The outcome of this research network, involving 31 research institutes in 17 countries, was summarized by Pereira et al. (2013a). Since then, the fields of applying nutritional, hormonal, and semiochemical treatments, as well as the work on optimizing pre-release conditions and operations that improve male quality, have continued to evolve. This chapter summarizes the current status in improving the performance of sterile males after mass-rearing.

2. NUTRITIONAL SUPPLEMENTS

In a majority of tephritid fruit flies, both males and females are anautogenous, emerging as adults with mostly undeveloped gonads. Both sexes rely on foraging during early adult life to ingest carbohydrates to fuel metabolic activities, as well as nitrogenous compounds for gonadal and accessory gland development and pheromone production (Epsky and Heath 1993; Hendrichs and Prokopy 1994). Thus, the survival and reproductive success of males in most tephritid species of economic importance is linked to the access to these nutrients. Nevertheless, several fly emergence and release facilities still do not include nitrogenous compounds in the pre-release diet of sterile males. The sterile males are generally offered a pre-release diet of only sugar or highly concentrated sucrose, presented in agar blocks, which are also a source of water (USDA/APHIS 2009; FAO/IAEA 2017).

The results on assessing nutritional supplements in the last 20 years are described thematically in more detail, together with relevant references, in Table 1. Studies on four genera of tephritids (*Anastrepha*, *Bactrocera*, *Ceratitis* and *Zeugodacus*) indicate that providing yeast hydrolysate to males, as part of the pre-release diet in the days following emergence, can enhance male sexual performance and reduce wild-female remating, although the optimal dosage and most practical form of presentation in an operational context still needs to be refined further (Kaspi and Yuval 2000; Aluja et al. 2001; Pérez-Staples et al. 2007; Haq et al. 2010a, 2014a). The implementation of such a strategy depends on a specific understanding of the nutritional needs of males (while feeding on nitrogenous sources in nature), trade-offs between sexual performance and survival, and operational costs and benefits. Another benefit of providing a protein-enriched pre-release diet is that sterile males will be much less attracted to nitrogenous baits in sprays or traps (such as Biolure or CeraTrap) that are increasingly being used for monitoring wild females when releasing only sterile males (Vargas et al. 2002; Maor et al. 2004; Vargas and Prokopy 2006; San Andrés et al. 2009).

Table 1. Summary of results, by thematic subject, of research conducted on the incorporation of nutritional supplements into pre-release diets and male performance of tephritids of the genera *Anastrepha*, *Bactrocera*, *Ceratitis*, and *Zeugodacus* (modified after Pereira et al. 2013a, reproduced with permission)

Subject addressed	Summary of results
Natural food	<ul style="list-style-type: none"> • Males feeding on orange (<i>Citrus sinensis</i> (L.) Osbeck) or mango (<i>Mangifera indica</i> L.) exhibit modified pheromone composition and improved sexual performance in <i>Anastrepha ludens</i> (Loew) and <i>Anastrepha obliqua</i> (Macquart) (Liedo et al. 2013; Utgés et al. 2013). • Male pheromone production is quantitatively affected by natural food in <i>A. ludens</i> and <i>A. obliqua</i> (Liedo et al. 2013). • Wild <i>Bactrocera tryoni</i> (Froggatt) males feeding on natural sources of food mature much slower than wild males with access to yeast hydrolysate; effects of diet are far less pronounced in mass-reared males (Weldon and Taylor 2011).
Dietary supplements	<ul style="list-style-type: none"> • Adding yeast hydrolysate (protein) to the pre-release diet significantly improves male sexual performance in <i>Anastrepha fraterculus</i> (Wiedemann), <i>A. ludens</i>, <i>A. obliqua</i>, <i>Anastrepha serpentina</i> (Wiedemann), <i>Anastrepha striata</i> Schiner, <i>Anastrepha suspensa</i> (Loew), <i>Bactrocera correcta</i> (Bezzi), <i>Bactrocera dorsalis</i> (Hendel), <i>B. tryoni</i>, <i>Bactrocera zonata</i> (Saunders), <i>Ceratitis capitata</i> (Wiedemann), <i>Ceratitis quilicii</i> De Meyer, Mwatawala, and Virgilio, and <i>Zeugodacus cucurbitae</i> (Coquillett) (Taylor and Yuval 1999; Aluja et al. 2001; Shelly et al. 2005; Pérez-Staples et al. 2007, 2008b, 2009; Yuval et al. 2007; Pereira et al. 2009, 2010a, b, 2011; Haq et al. 2010a, b, 2013; Haq and Hendrichs 2013; Liedo et al. 2013; Liendo et al. 2013; Orankanok et al. 2013; Quilici et al. 2013; Ndzana et al. 2016; Shelly 2017a). • Adding protein to the pre-release diet improves male weight and body protein content in <i>A. suspensa</i>, <i>B. dorsalis</i>, and <i>Z. cucurbitae</i> (Haq et al. 2010c; Pereira et al. 2011; Reyes-Hernández et al. 2019). It also enhances male sexual organ development in <i>B. dorsalis</i> (Reyes-Hernández et al. 2019) and <i>B. tryoni</i> (Pérez-Staples et al. 2011). • Some studies in <i>C. capitata</i> found no difference between sugar-only fed males and yeast hydrolysate-fed males in terms of mating success (Shelly and McInnis 2003; Shelly et al. 2003, 2006a), survival rate (Shelly and McInnis 2003), and dispersal (Shelly and Edu 2008a).
Ratio of protein to carbohydrates	<ul style="list-style-type: none"> • Protein in dry diets early in adult life contributes to male sexual performance, but the ratio of protein to carbohydrates affects survival in a dose-dependent manner in <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, and <i>B. tryoni</i> (Prabhu et al. 2008; Gómez et al. 2013; Liedo et al. 2013; Liendo et al. 2013; Utgés et al. 2013). • As little as 4–10% of protein content in the pre-release diet is sufficient to significantly enhance male sexual performance in <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, and <i>B. tryoni</i> (Pérez-Staples et al. 2008a; Pereira et al. 2011; Gómez et al. 2013; Liedo et al. 2013; Liendo et al. 2013).
Optimal formulation/delivery system	<ul style="list-style-type: none"> • A formulation based on soy whey protein and incorporating methoprene has been developed and tested for <i>A. suspensa</i> (Teal et al. 2013). • Concurrently, a commercial product "Mubarqui" containing protein has been introduced in a dry formulation into release programmes for <i>A. ludens</i> and <i>A. obliqua</i> (Gómez and Teal 2010; Gómez et al. 2013).
Dietary effects on remating inhibition	<ul style="list-style-type: none"> • Feeding on a nitrogen-rich food significantly improves sterile male ability to inhibit female remating in <i>A. fraterculus</i> (Abraham et al. 2011, 2012, but see Abraham et al. 2013), <i>Bactrocera carambolae</i> Drew and Hancock, <i>B. tryoni</i>, <i>C. capitata</i>, and <i>Z. cucurbitae</i> (Pérez-Staples et al. 2008a; Gavriel et al. 2009; Haq et al. 2014a).

Table 1. Continued

Subject addressed	Summary of results
Interaction between diet, dispersal, and survival	<ul style="list-style-type: none"> • Protein-rich diets are related to a lower resistance to starvation in <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, <i>C. capitata</i>, and <i>B. tryoni</i> (Levy et al. 2005; Utgés et al. 2013; Reynolds et al. 2014; Juárez et al. 2019). • Adult diets that contain protein are related to lower recapture rates in protein-baited traps, and to reduced mobility and longevity in sterile <i>A. ludens</i>, <i>A. obliqua</i> (but not in <i>C. capitata</i>) males (Maor et al. 2004; San Andrés et al. 2009; Utgés et al. 2013). • Protein-rich post-teneral diets do not adversely affect the survival and movement in the field of sterile <i>B. tryoni</i> and <i>C. capitata</i> males (Yuval et al. 2007; Gavriel et al. 2010; Taylor et al. 2013b). • In the laboratory, protein-fed males live significantly longer than only sugar-fed males of <i>B. tryoni</i> and <i>Z. cucurbitae</i> (Haq et al. 2010b; Taylor et al. 2013b). The same was found for <i>A. ludens</i> and <i>A. obliqua</i> when the sugar:yeast ratio is 9:1 or 24:1, but not with a 3:1 ratio (Liedo et al. 2013). • In the laboratory, <i>B. tryoni</i> fed a 3:1 ratio of sugar:yeast for two days following emergence are more vulnerable to starvation if food is unavailable in the following days than if the flies are fed only sugar (Taylor et al. 2013a).
Contribution of micro-organisms	<ul style="list-style-type: none"> • Supplementing post-teneral diets with symbiotic bacteria reduces mass-rearing costs and increases mating success in fruit fly pests in SIT application (Cáceres et al. 2019; Noman et al. 2020; Raza et al. 2020; Augustinos et al., this volume). • The significant contribution of micro-organisms to the fitness of non-sterile <i>Bactrocera oleae</i> (Rossi) and <i>C. capitata</i> has been established (Behar et al. 2008a; Ben-Yosef et al. 2008a, b, 2010, 2014, 2015; Yuval et al. 2013; Stathopoulou et al. 2021). In <i>B. oleae</i>, the survival of males feeding on a probiotic diet is reduced compared with those fed on a sugar-only diet (Sacchetti et al. 2014). • Presence of commensal bacteria affect mate-selection behaviour of <i>B. dorsalis</i> (Damodaram et al. 2016). • Manipulating the microflora of sterile males by supplementing probiotic bacteria in adult diets can improve male sexual performance in <i>C. capitata</i> (Niyazi et al. 2004; Behar et al. 2005, 2008a, b, c; Ben-Ami et al. 2010; Gavriel et al. 2011; Hamden et al. 2013; Kyritsis et al. 2019), and in <i>A. obliqua</i>, <i>B. dorsalis</i>, and <i>Z. cucurbitae</i> (Rull et al. 2015; Yao et al. 2017; Cai et al. 2018; Stathopoulou et al. 2021).
Irradiation effects mitigated by dietary supplements	<ul style="list-style-type: none"> • Feeding with live bacteria can restore ecological fitness of males that were sterilized through irradiation in <i>B. dorsalis</i> (Cai et al. 2018). • Diet and irradiation affect micro-organism diversity in <i>B. tryoni</i> (Woruba et al. 2019). • Microflora diversity in laboratory-reared and irradiated males is low when compared with wild <i>C. capitata</i> males (Ben-Ami et al. 2010). • Irradiation significantly diminishes the ability of <i>B. tryoni</i> flies to tolerate protein deprivation (Pérez-Staples et al. 2007; Taylor et al. 2013b).

Studies on the nutritional requirements of males of several tephritid species and their effect on sexual success (involving various species targeted by the SIT) have reached different levels of understanding. For example, studies in nature on *C. capitata* provide evidence that wild males feed on sources of protein (Hendrichs and Hendrichs 1990; Hendrichs et al. 1991). However, while providing yeast hydrolysates or other sources of protein together with sugar to males generally improved mating success in *Anastrepha*, *Bactrocera*, and *Zeugodacus* (Table 1), in *Ceratitis* there is a variance in results reflecting the influence of other factors.

Different effects of protein-enriched diets have been found, in some cases reducing male survival and dispersal; the adverse effect was reduced when the protein ratio in the diet was lowered, but there are no clear procedural guidelines available currently for this and other target species (Blay and Yuval 1997; Shelly and Kennelly 2003; Shelly and McInnis 2003; Prabhu et al. 2008; FAO/IAEA 2017).

Moreover, there is evidence that the integrity of micro-organisms in the male gut contribute to fly health, foraging behaviour, and sexual performance (Yuval et al. 2013; Akami et al. 2019; Deutscher et al. 2019; Juárez et al. 2019; Augustinos et al., this volume). Adult diets enriched with different probiotic bacteria strains showed in irradiated *C. capitata* males a positive effect on the mating competitiveness, mating latency time, sexual calling or longevity (Niyazi et al. 2004; Behar et al. 2008b; Ben Ami et al. 2010; Gavriel et al. 2011).

Nevertheless, despite the increasing scientific evidence of the benefits of nutritional supplements, currently, possibly out of convenience, sterile males are often offered only a pre-release diet of highly concentrated sucrose, normally presented in an agar block, but without any protein addition. The formulation and testing of optimal pre-release diets, containing sugar, protein and probiotic bacteria (and possibly other ingredients) in proportions that will result in enhanced sterile male performance in the field, is still not fully understood and accepted by many operational programme managers, and remains to be implemented. However, the Moscamed programme in Mexico uses the Mubarqui adult diet for *C. capitata*, which contains proteins from diverse plant seeds (Gómez et al. 2013), and the Moscafrut programme in Mexico releases sterile *A. ludens* and *A. obliqua* flies fed with a 24:1 sugar:yeast adult diet. Managers should evaluate this matter in relation to their fruit fly species and decide on the most appropriate pre-release adult diet and feeding regime for sterile males.

3. HORMONAL TREATMENTS

Age is a significant factor affecting sexual signalling and reproduction in numerous tephritid species (Liedo et al. 2002; Pereira et al. 2013a). Being anautogenous, tephritid fruit fly pests in nature require a considerable period (over a week in *Ceratitidis* spp.; 2–3 weeks in *Anastrepha* spp., *Bactrocera* spp., and *Zeugodacus* spp.) under adequate nutritional conditions to reach sexual maturation (Teal et al. 2013). Even though mass-rearing inadvertently selects for accelerated sexual maturation, the discrepancy between the standard holding periods at fly emergence and release facilities, and the time to reach sexual maturation, poses a significant problem for SIT programmes. After release but before reaching full sexual maturation, sterile males suffer significant losses due to predation and other causes, resulting in far fewer males surviving to maturity and copulation (Hendrichs and Hendrichs 1998; Rao et al. 2013; González-López et al. 2016).

Holding sterile males for more days also increases the management costs, and additional investment is required to expand the infrastructure at fly emergence and release facilities (USDA/APHIS 2009). In addition, in those programmes where sterile males are released together with sterile females, longer holding times are not advisable because males may start mating before release, thereby transferring their

limited sperm to sterile females. Clearly, the development of cost-effective methods to accelerate sexual maturity in released flies would have a significant positive impact on the efficacy of the SIT (Teal et al. 2000).

Research on several tropical *Anastrepha* species shows that juvenile hormone is a critical hormone, regulating sexual maturity and sexual signalling in these species (Table 2). Furthermore, application of juvenile hormone or the analogues, methoprene or fenoxycarb, can accelerate reproductive development and sexual signalling in sterile males of some species (Pereira et al. 2013a, b; Teal et al. 2013). Significant progress has been made in evaluating hormonal treatments using methoprene to accelerate reproductive development and, in some cases, improving further male sexual performance. It was found in many species (but not all) that sterile males become sexually mature significantly earlier when hormone exposure is included in pre-release holding protocols, but this optimum effect was only achieved when hormone treatment was coupled with a protein-enriched pre-release diet (Teal et al. 2013). Thus, irradiated males become sexually mature earlier and can be released earlier. This advantage is particularly important for SIT application against those species of *Anastrepha* spp., *Bactrocera* spp., and *Z. cucurbitae* that have long pre-copulatory periods. The improvement in male sexual performance by applying hormonal treatments was extensively studied in *Anastrepha* spp., *B. dorsalis*, *B. tryoni*, *Z. cucurbitae*, and *C. capitata*, although in these last two species no benefits were found. In addition, considerable progress was made in developing delivery systems to treat large numbers of flies with methoprene in operational programmes. The results on hormonal treatment assessments are described thematically in more detail, together with the relevant references, in Table 2.

Table 2. Summary of results, by thematic subject, of research conducted on the effects of pre-release application of hormonal treatments on male performance of tephritids of the genera *Anastrepha*, *Bactrocera*, *Ceratitis*, and *Zeugodacus* (modified after Pereira et al. 2013a, reproduced with permission)

Subject addressed	Summary of results
Age of sexual maturation of males determined	<ul style="list-style-type: none"> Age at which laboratory-reared and/or wild males of <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, <i>A. serpentina</i>, <i>A. striata</i>, <i>A. suspensa</i>, <i>B. correcta</i>, <i>B. dorsalis</i>, <i>B. tryoni</i>, <i>C. capitata</i>, <i>C. quilicii</i>, and <i>Z. cucurbitae</i> become sexually mature has been determined (Teal and Gómez-Simuta 2002; Aluja et al. 2009; Pereira et al. 2009; Gómez et al. 2013; Obra and Resilva 2013; Orankanok et al. 2013; Pereira et al. 2013b; Quilici et al. 2013; Segura et al. 2013; Sookar et al. 2013; Adnan 2019).
Improvement in reproductive maturation and sexual performance	<ul style="list-style-type: none"> Hormonal treatment has positive effects on accelerating reproductive development and improving sexual performance in <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, <i>A. serpentina</i>, <i>A. striata</i>, <i>A. suspensa</i>, <i>B. tryoni</i>, and <i>Z. cucurbitae</i> (Aluja et al. 2009; Pereira et al. 2009; Segura et al. 2009; Haq et al. 2010a, 2013; Gómez et al. 2013; Collins et al. 2014; Gomez-Simuta et al. 2017; Bachmann et al. 2017; Adnan et al. 2020a). In <i>A. fraterculus</i>, the accessory glands of males treated with methoprene are less effective at inhibiting female remating than males not treated with methoprene (Abraham et al. 2012). There is no effect of hormonal treatment on either <i>B. dorsalis</i> or <i>C. capitata</i> (Faria et al. 2008; Shelly et al. 2009).

Table 2. Continued

Subject addressed	Summary of results
Interaction of protein and hormone treatments on sexual performance	<ul style="list-style-type: none"> • Males treated with hormone and feeding on a protein-rich diet effectively compete with mature wild males, but at significantly earlier ages for <i>A. fraterculus</i>, <i>A. ludens</i>, and <i>A. suspensa</i> (Gómez and Teal 2010; Liendo et al. 2013; Pereira et al. 2013b; Gómez et al. 2013). • Males of <i>Z. cucurbitae</i> treated with methoprene and with access to a protein-rich diet have an accelerated maturation and improved mating performance (Haq et al. 2010a).
Interaction of irradiation and hormone treatments on sexual maturity	<ul style="list-style-type: none"> • Interaction of irradiation and hormone treatments has no negative effects on male acceleration of reproductive maturity in <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, <i>A. suspensa</i>, and <i>B. tryoni</i> (Teal et al. 2007; Segura et al. 2013; Adnan et al. 2019). • Females mated with 6-day-old, methoprene-treated <i>A. fraterculus</i> males remated more and sooner than females mated with naturally matured males, either sterile or wild (Abraham et al. 2013).
Optimal dose to accelerate maturation	<ul style="list-style-type: none"> • Optimal hormone treatment dose has been determined using topical application for <i>A. fraterculus</i>, <i>A. ludens</i>, <i>A. obliqua</i>, <i>A. suspensa</i>, and <i>Z. cucurbitae</i> (Teal et al. 2007; Pereira et al. 2009; Segura et al. 2013; Haq et al. 2010a).
Response in males and females to hormone treatments	<ul style="list-style-type: none"> • A differential response in males and females to methoprene exposure is found in <i>A. fraterculus</i> and <i>B. tryoni</i>; it can act as a physiological sexing system, minimizing matings between sterile males and sterile females (Segura et al. 2009; Liendo et al. 2013; Adnan et al. 2019). • No differential response to methoprene exposure has been found in <i>A. ludens</i> males and females (Pereira et al. 2013b).
Development of delivery systems other than topical application	<ul style="list-style-type: none"> • Agar-based diet with 5–10% protein along with 0.05% methoprene, tested with <i>A. ludens</i> and <i>A. suspensa</i>, produces large amounts of waste and is not cost-effective (Pereira et al. 2013b; Teal et al. 2013). • Pupal dipping in an acetone bath containing 0.05% methoprene, tested with <i>A. fraterculus</i>, <i>A. ludens</i>, and <i>A. suspensa</i>, does not impact emergence or survival and accelerates reproductive development, although it can cause humidity problems when handling large volumes of pupae (Liendo et al. 2013; Pereira et al. 2013b; Segura et al. 2013). • Pupal dipping is impractical due to health risks, safety, and disposal issues associated with the use of acetone (Pereira et al. 2013a). • A dry sugar–protein pre-release diet containing 0.0015% methoprene for <i>A. ludens</i> and <i>A. obliqua</i> is practical and effective in accelerating development when applied in operations of fly emergence and release facilities (Gómez et al. 2013). • Dietary treatment with mosquito larvicide containing methoprene in a dry 3:1 mix of sugar to yeast is effective in <i>B. tryoni</i> (Adnan 2019; Adnan et al. 2019, 2020a).
Efficient large-scale methods for incorporation of JH analogues	<ul style="list-style-type: none"> • The dry sugar–protein–methoprene diet method was assessed in a 3500-ha pilot area for <i>A. ludens</i> in Mexico (Gómez et al. 2013). • The hormone and protein delivery system has tested positive in the “Moscafrut” action programme to control <i>A. ludens</i> fruit flies in Mexico (Gómez et al. 2013).

4. SEMIOCHEMICAL TREATMENTS

Males of most *Anastrepha* spp., *Bactrocera* spp., *Ceratitis* spp., and *Zeugodacus* spp. are attracted to natural compounds known as semiochemicals (Cunningham 1989; Kumaran et al. 2013; Segura et al. 2018). Some species are known to sequester these chemicals from plants for use in pheromone synthesis; for example, ingestion of methyl eugenol (ME) from natural sources by males of *B. dorsalis* and *B. carambolae* results in the storage of metabolites in the rectal gland and their subsequent release as part of the pheromone during fanning performed in courtship (Tan and Nishida 1996). Providing sterile *B. dorsalis* males with a source of ME to feed on before release could increase their mating competitiveness (Shelly et al. 2005). ME feeding may also confer a survival advantage because metabolites appear to act as a very potent allomone to deter vertebrate predators (Wee and Tan 2001).

In the case of *C. capitata*, exposure to ginger (*Zingiber officinale* Roscoe) root oil (GRO) or citrus oils (by contact or vapour) enhances considerably the mating competitiveness of wild or mass-reared males (Katsoyannos et al. 1997; Papadopoulos et al. 2001; Shelly 2001).

Exposure to guava (*Psidium guajava* L.) fruit volatiles increased the mating success of both wild and laboratory *A. fraterculus* males relative to non-exposed males (Vera et al. 2013). Also, citrus and grapefruit volatiles have potential to enhance the mating performance of male *A. ludens* and *A. fraterculus* (Bachmann et al. 2015; Morató et al. 2015). However, the attraction and mating enhancing effects of the semiochemicals so far found for *Anastrepha* effects are not as dramatic as seen with *Bactrocera*, *Ceratitis* and *Zeugodacus*, and suggest that similar compounds may exist in the *Anastrepha* ecological sphere and await discovery (Segura et al. 2018).

The research carried out in this area has helped to understand these phenomena, to extend them to other species, and to transfer and validate them under the large-scale conditions of action SIT programmes. Results obtained recently confirmed the compounds that have the potential to improve sterile male performance of target fruit flies in the field, and some additional ones have been identified. These chemicals are the following:

- Methyl eugenol (ME) is found in more than 450 plant species (Tan and Nishida 2012); exposure to holy basil (*Ocimum tenuiflorum* L.) and sweet basil (*Ocimum basilicum* L.) oils, as well as ripe fruits of tropical almond (*Terminalia catappa* L.), improve the mating competitiveness of several *Bactrocera* species, including *B. correcta*, *B. dorsalis*, and *B. zonata* males (Cunningham 1989; Tan and Nishida 1996; Quilici et al. 2004; Shelly et al. 2005; Shelly and Edu 2007; Obra and Resilva 2013; Orankanok et al. 2013).
- Cuelure (CUE) has potential to enhance the performance of *B. tryoni* and *Z. cucurbitae* males (Weldon et al. 2008; Shelly 2019).
- Raspberry ketone (RK) and/or zingerone (ZG) have potential to enhance attraction, maturation, and sexual performance of *B. tryoni* and *Z. cucurbitae* males (Khoo and Tan 2000; Akter et al. 2017b; Akter and Taylor 2018; Shelly 2019); however, Fezza and Shelly (2018) did not find that access to a RK-supplemented diet at an early age accelerated sexual maturation.

- Ginger root oil (GRO) and α -copaene enhance male performance of *C. capitata* and *C. quilicii* (Shelly 2001; Shelly et al. 2007a; Quilici et al. 2013).
- Manuka oil (rich in α -copaene) from the manuka tree (*Leptospermum scoparium* Forst and Forst) of New Zealand improves *C. capitata* male mating (Shelly et al. 2008c).
- Orange oils (OO) and other citrus oils enhance male performance of *C. capitata* and *C. quilicii*, and potentially *A. fraterculus* (Shelly 2001; Shelly et al. 2007a; Quilici et al. 2013; Vera et al. 2013).
- Citrus, grapefruit, and guava fruit volatiles have potential to enhance the mating performance of male *A. ludens* and *A. fraterculus* (Vera et al. 2013; Bachmann et al. 2015; Morató et al. 2015).

The results of assessing semiochemical treatments, together with the relevant references, are described in more detail in Table 3.

Methodologies for exposing large numbers of *C. capitata* males through GRO or citrus-oil aromatherapy on a large scale in adult-holding rooms at fly emergence and release facilities have been developed (Shelly et al. 2007c, 2008a). They are now applied routinely in a very cost-effective manner in on-going SIT programmes in Australia, Croatia, Guatemala, Israel, Mexico, Spain, and the USA, resulting in a significant improvement in mating performance of the released sterile males.

It is envisaged that providing sterile *Bactrocera* spp. males with a source of ME to feed on before release will place them on at least an even “playing field” against wild males, thereby potentially reducing the number or frequency of sterile males released (Barclay et al. 2014; Vargas et al. 2014). However, the common methods for holding sterile males in fly emergence and release facilities do not enable the feeding of ME to millions of adult flies after their emergence. Additionally, exposure to ME must be brief in view of ME toxicity when access is unlimited. In response, Tan and Tan (2013) designed an automated ME machine prototype for briefly feeding ME to sterile males on a belt impregnated with ME after which they are brushed off and collected. However, this approach is not really suitable for industrial processing of millions of sterile males. Considering this scenario at fly emergence and release facilities, Haq et al. (2014b, 2015, 2018) demonstrated that ME application by aromatherapy also enhanced the mating success of males of *B. carambolae* and *B. dorsalis*, finding that ME aromatherapy produced a mating boost as early as one day after exposure. This alternative method appears to have merit for adoption but needs to be evaluated at larger scales in fly emergence and release facilities.

Another advantage of exposure to ME is that it reduces significantly the response of sterile males to ME in male annihilation treatments, thus potentially enabling the simultaneous application of the SIT and male annihilation (Barclay et al. 2014; Vargas et al. 2014). Similarly, pre-release exposure to RK has been found to diminish the subsequent response of *B. tryoni* to CUE sources (Akter et al. 2017a).

Table 3. Summary of results, by thematic subject, of research conducted on the effects of pre-release application of semiochemical treatments on male performance of tephritids of the genera *Anastrepha*, *Bactrocera*, *Ceratitis*, and *Zeugodacus*. Abbreviations explained in the text (modified after Pereira et al. 2013a, reproduced with permission)

Subject addressed	Summary of results
Search for semiochemical compounds that affect <i>Anastrepha</i> spp. male sexual performance	<ul style="list-style-type: none"> • Grapefruit volatiles increase male mating competitiveness in <i>A. ludens</i> (Morató et al. 2015). • Male mating performance of <i>A. fraterculus</i> is positively affected by exposure to guava volatiles (Vera et al. 2013; Bachmann et al. 2015). • Exposure to lemon fruit volatiles enhances male mating performance in <i>A. fraterculus</i>, but there are also some detrimental effects (Vera et al. 2013). GRO exposure also increases male mating success in <i>A. serpentina</i> (Flores et al. 2011).
Assessment of various semiochemicals in enhancing <i>Bactrocera</i> spp. male performance	<ul style="list-style-type: none"> • Exposure to basil oil (containing ME) has a positive effect on sterile male mating performance of <i>B. dorsalis</i> (Obra and Resilva 2013). • Exposure to ripened tropical almond fruit (containing ME) enhances <i>B. dorsalis</i> male mating success (Shelly and Edu 2007). • Exposure to ME significantly improves sterile male mating performance of <i>B. correcta</i> and <i>B. dorsalis</i> (Shelly et al. 1996, 2005, 2008b, 2009, 2010a; Hee and Tan 1998; Shelly and Edu 2008b; Ji et al. 2013; Orankanok et al. 2013). • Some negative effects on the male ejaculate have been found in <i>B. dorsalis</i> (Reyes-Hernández et al. 2019). • ME-mediated male mating enhancement also found in <i>Bactrocera cacuminata</i> (Hering) (Raghu and Clarke 2003), <i>B. carambolae</i> (Wee et al. 2007), and <i>Bactrocera umbrosa</i> (F.) (Wee et al. 2018). • Exposure to commercial ME, and interaction with post-teneral adult diet, increases male mating performance of <i>B. correcta</i> and <i>B. dorsalis</i>. For both species, ME-treated sterile males are being released in operational SIT programmes in Thailand (Orankanok et al. 2013). • Sterile males, feeding on commercial ME prior to release, induce higher egg sterility in wild populations than control sterile males denied ME (McInnis et al. 2011). • Exposure to ME does not affect dispersal or survival of sterile flies of <i>B. correcta</i> and <i>B. dorsalis</i> (Orankanok et al. 2013).
Assessment of various semiochemicals in enhancing male performance of other <i>Bactrocera</i> spp.	<ul style="list-style-type: none"> • Feeding on CUE increases <i>Z. cucurbitae</i> mating success (Shelly and Nishimoto 2017). Feeding on RK enhances male mating performance of <i>Z. cucurbitae</i>, but only for one day post-feeding (Shelly 2000, 2019). However, exposure to zingerone has no effect on <i>Z. cucurbitae</i> male mating competitiveness (Shelly 2017b) or a positive effect only one day after exposure (Inskeep et al. 2019). • Exposure to CUE and RK increases the mating competitiveness of <i>B. tryoni</i>, but young males fail to respond to CUE, OO, and GRO (Weldon et al. 2008; Kumaran et al. 2014a; Akter et al. 2017b). • <i>B. tryoni</i> feeding on yeast hydrolysate enhances attraction to CUE (Weldon et al. 2008), and on CUE increases male signalling, female attraction, and male mating success (Kumaran et al. 2013, 2014b). • <i>B. tryoni</i> feeding on RK (or caffeine supplements) promotes male early sexual maturation, greater mating success and increased multiple mating, without affecting remating receptivity in females (Akter et al. 2017b; Akter and Taylor 2018; Khan et al. 2019; Adnan et al. 2020b; GFV 2020). • Exposing sexually mature adult <i>B. oleae</i> to the aroma of α-pinene significantly increases the mating performance over non-exposed males and females (Gerofotis et al. 2013; Kokkari et al. 2017).

Table 3. Continued

Subject addressed	Summary of results
Optimal age and feeding dose for <i>Bactrocera</i> spp. male flies	<ul style="list-style-type: none"> • For <i>B. dorsalis</i>, Obara and Resilva (2013) determined the optimal age and duration of ME exposure, and the diurnal pattern of ME feeding. • Optimal age of ME exposure, in combination with nutritional supplements, has been determined for sterile <i>B. correcta</i> and <i>B. dorsalis</i> males (Orankanok et al. 2013).
Effect of semiochemical treatments on <i>C. capitata</i> sexual signalling (pheromone calling)	<ul style="list-style-type: none"> • GRO-exposed males exhibit higher rates of sexual signalling compared with unexposed <i>C. capitata</i> males (Shelly 2001, 2007a, 2008, 2010b; Papadopoulos et al. 2006), although the effect found by Briceño et al. (2007) was lower. • During sexual courtship GRO-exposed sterile males exhibit the same wing-beat duration as wild males, and this is shorter than for unexposed sterile males (Morelli et al. 2013). • Exposure to OO and their components increases sexual signalling and mating success of <i>C. capitata</i> males (Shelly et al. 2004, 2006b; Kouloussis et al. 2013); this effect is more pronounced in protein-fed males, independent of age (Katsoyannos et al. 2004; Papadopoulos et al. 2006; Kouloussis et al. 2013, 2017). • Exposure to volatiles of leaves of the tea tree increases mating competitiveness of sterile <i>C. capitata</i> males (Shelly and Epsky 2015).
Identification of active compounds other than GRO, and their effects on sexual performance of <i>C. capitata</i>	<ul style="list-style-type: none"> • Exposure to oranges and grapefruits (<i>Citrus paradisi</i> Macfad) improves mating success of male <i>C. capitata</i> (Papadopoulos et al. 2001; Shelly 2009). • Exposure to fruits of five citrus species shows that sweet oranges confer the highest increase in male mating performance of <i>C. capitata</i>. Exposure to commercial citrus oils is similarly effective (Kouloussis et al. 2013). • There is a positive effect of OO components (limonene, β-myrcene, and linalool) on protein/sugar-feeding males, but not on sugar-only-feeding males (Kouloussis et al. 2013). • A mixture of limonene, β-myrcene, linalool, α-pinene, and geraniol (1:1:1:1:1) is very effective for both wild and sterile <i>C. capitata</i> male sexual performance (Kouloussis et al. 2013).
Effects of GRO and OO exposure to <i>C. quilibii</i>	<ul style="list-style-type: none"> • Exposure of sugar-only-fed <i>C. quilibii</i> males to GRO and OO increases their mating performance, but only GRO increases the mating performance of protein/sugar-fed males (Quilibi et al. 2013).
Effect of semiochemical treatments on <i>C. capitata</i> female remating frequency	<ul style="list-style-type: none"> • The remating frequency of <i>C. capitata</i> females mated with sterile males treated with 0.1 ml/m³ GRO is similar to that of females mated with wild males and lower than that of females mated with sterile unexposed males (Morelli et al. 2013).
Effect of semiochemical treatments on <i>C. capitata</i> dispersal and survival	<ul style="list-style-type: none"> • Field dispersal and survival rates of GRO-exposed sterile males are similar to those of unexposed <i>C. capitata</i> males (San Andrés et al. 2009; Paranhos et al. 2010; Juan-Blasco et al. 2013).
Determining the dose of semiochemical exposure for <i>C. capitata</i>	<ul style="list-style-type: none"> • OO compounds show positive effects in increasing <i>C. capitata</i> male performance (Kouloussis et al. 2013). The optimal dose of OO exposure was determined for both protein-sugar and sugar-only <i>C. capitata</i> adult diets. There are positive effects of OO for a protein-sugar diet, but no effects for a sugar-only diet (Kouloussis et al. 2013).
Optimal delivery system for large-scale semiochemical application in <i>Ceratitidis</i> spp. SIT programmes	<ul style="list-style-type: none"> • The cost-effective GRO exposure to sterile males has been established, and is in operation in several <i>C. capitata</i> SIT programmes (Shelly et al. 2007b, c, 2008a; Paranhos et al. 2008, 2013; Juan-Blasco et al. 2013; Silva et al. 2013; Steiner et al. 2013).

5. CONCLUSIONS

Extensive laboratory and semi-field tests indicate that pre-release diets nutritionally enriched with some yeast hydrolysate improve sterile male performance; they can result in increased SIT efficiency for a majority of fruit fly species tested, and thus can be adopted by programme managers, although some confounding factors still need to be better understood for some target species, as well as the optimal protein to carbohydrate ratios and physical form of diet provision. Also, a better understanding of the effect of nutrition during the larval stage on the nutritional requirements of the adults is needed (Aluja et al. 2009).

In addition, hormone therapy coupled with feeding protein to adults generally accelerates male maturation and sexual performance among *Anastrepha* species and some *Bactrocera* species, with no negative side effects on survival. Furthermore, in some species, methoprene treatment has increased male mating and competitiveness. However, negative effects have been recorded on female remating in *A. fraterculus*, and on sperm storage and male ejaculate in *B. dorsalis*.

Accelerated male maturation and improved sexual performance can represent significant cost savings associated with infrastructure investment, reduced holding periods for sterile males prior to release, and fewer males dying in the field before reaching sexual maturity. Validation in a few fruit fly species has resulted in some of these major breakthroughs being incorporated into several action programmes, although others, for technical or practical reasons, have been reluctant to incorporate these innovations into their operations.

The development of cost-effective semiochemical treatments that improve sterile male sexual signalling and attractiveness to wild females, and thus overall mating performance, is highly desirable. Implementation of aromatherapy has significantly increased the effectiveness of on-going programmes integrating the SIT against *C. capitata*. On the other hand, while the incorporation of semiochemical treatments for *Bactrocera* species appears to be feasible, it still remains to be implemented operationally; major benefits result from the simultaneous field application of male annihilation to eliminate wild males. Thus, incorporating an ME pre-release treatment has considerable potential to significantly increase sterile to wild male overflooding ratios (Barclay et al. 2014; Vargas et al. 2014; Hendrichs and Robinson, this volume). For some *Anastrepha* species and other species such as *B. oleae* and *C. quilibii*, potentially similar semiochemicals and fruit volatiles probably exist, and more research funding should be allocated to find and investigate them (Segura et al. 2018).

Also critical for sterile male performance are the processes to which sterile males are subjected at fly emergence and release facilities (USDA/APHIS 2009; FAO/IAEA 2017; Dowell et al., this volume), the environmental conditions in which they are held, and their interaction with exposure to hormonal, nutritional and/or semiochemical treatments (Pereira et al. 2013a). In addition, holding densities (Díaz-Fleischer et al. 2009; Andress et al. 2015), the timing of providing treatments in relation to fly age and sexual maturation, and holding periods are also very relevant. For example, results indicate that in *C. capitata* sexing strains, the release of older males, closer to sexual maturity, is preferable; this results in more flies reaching mating age in the field (McInnis et al. 2013). Species with longer pre-copulatory

maturation periods, such as *Anastrepha* spp. and *Bactrocera* spp., benefit more from combined nutritional and hormonal treatments to reduce long pre-release holding periods (Gómez et al. 2013; Pereira et al. 2013a).

Assessing the interaction of different environmental and holding conditions, combined with hormonal, nutritional, and/or semiochemical treatments, on sterile fly performance has resulted in the routine application of GRO under large-scale holding conditions in operational *C. capitata* SIT programmes (USDA/APHIS 2009; FAO/IAEA 2017). Also, a hormone-and-protein-delivery system has been developed for the SIT programmes suppressing *A. ludens* fruit fly populations in Mexico (Gómez et al. 2013). Furthermore, the effects on fly performance (in terms of dispersal and recapture) of different sterile fly shipping and release systems, and their interaction with hormonal, nutritional, and/or semiochemical treatments, have been assessed for some aerial- and ground-release systems for *C. capitata* (Paranhos et al. 2010, 2013; Silva et al. 2013). Finally, parameters of sterile-fly quality, treatments, space, labour, and economics were assessed and compared at eight fly emergence and release facilities in Guatemala, Mexico, and the USA (USDA/APHIS 2009); there is a good baseline on which to improve.

Knowledge gained so far, and the practical procedures developed, are transferable, at least in part, to other insect pest species with management programmes that include an SIT component. Nevertheless, much remains unknown about the effects of manipulating the environmental holding conditions, either separately or in combination with nutritional, hormonal, and semiochemical treatments, on subsequent male quality in the field. Release methods, while operationally convenient, are not always optimal in terms of sterile male performance (Dowell et al., this volume). For example, the effect of chilling (to immobilize) collected flies for aerial release can at least temporarily affect sterile male flight ability and mating competitiveness, including pheromone quantity or quality (Andress et al. 2012, 2013; Shelly et al. 2013; Arredondo et al. 2016). Therefore, the effects and interactions of the different processes, treatments, and systems need to be further assessed and refined, tailoring them to the biology of each target fruit fly species.

6. REFERENCES

- Abraham, S., L. Goane, J. Cladera, and M. T. Vera. 2011. Effects of male nutrition on sperm storage and remating behavior in wild and laboratory *Anastrepha fraterculus* (Diptera: Tephritidae) females. *Journal of Insect Physiology* 57(11): 1501–1509. <https://doi.org/10.1016/j.jinsphys.2011.08.004>
- Abraham, S., J. Cladera, L. Goane, and M. T. Vera. 2012. Factors affecting *Anastrepha fraterculus* female receptivity modulation by accessory gland products. *Journal of Insect Physiology* 58: 1–6. <https://doi.org/10.1016/j.jinsphys.2011.08.007>
- Abraham, S., M. C. Liendo, F. Devescovi, P. A. Peralta, V. Yusef, J. Ruiz, J. L. Cladera, M. T. Vera, and D. F. Segura. 2013. Remating behavior in *Anastrepha fraterculus* (Diptera: Tephritidae) females is affected by male juvenile hormone analog treatment but not by male sterilization. *Bulletin of Entomological Research* 103(3): 310–317. <https://doi.org/10.1017/S0007485312000727>
- Adnan, S. M. 2019. Dietary methoprene and caffeine as pre-release supplements for Queensland fruit fly *Bactrocera tryoni* (Froggatt) sterile insect technique. PhD thesis, Macquarie University, Australia.
- Adnan, S. M., I. Farhana, J. R. Inskeep, P. Rempoulakis, and P. W. Taylor. 2019. Accelerated sexual maturation in methoprene-treated sterile and fertile male Queensland fruit flies (Diptera:

- Tephritidae), and mosquito larvicide as an economical and effective source of methoprene. *Journal of Economic Entomology* 112: 2842–2849. <https://doi.org/10.1093/jee/toz214>
- Adnan, S. M., I. Farhana, J. Inskeep, P. Rempoulakis, and P. W. Taylor. 2020a.** Dietary methoprene enhances sexual competitiveness of sterile male Queensland fruit flies in field cages. *Journal of Pest Science* 93: 477–489. <https://doi.org/10.1007/s10340-019-01170-0>
- Adnan, S. M., I. Farhana, S. J. Park, P. Rempoulakis, and P. W. Taylor. 2020b.** Caffeine as a promotor of sexual development in sterile Queensland fruit fly males. *Scientific Reports* 10: 14743. <https://doi.org/10.1038/s41598-020-71671-x>
- Akami, M., A. A. Andongma, C. Zhengzhong, J. Nan, K. Khaeso, E. Jurkevitch, C.-Y. Niu, and B. Yuval. 2019.** Intestinal bacteria modulate the foraging behavior of the oriental fruit fly *Bactrocera dorsalis* (Diptera: Tephritidae). *PLOS ONE* 14(1): e0210109. <https://doi.org/10.1371/journal.pone.0210109>
- Akter, H., and P. W. Taylor. 2018.** Sexual inhibition of female Queensland fruit flies mated by males treated with raspberry ketone supplements as immature adults. *Journal of Applied Entomology* 142: 380–387. <https://doi.org/10.1111/jen.12484>
- Akter, H., S. Adnan, R. Morelli, P. Rempoulakis, and P. W. Taylor. 2017a.** Suppression of culeure attraction in male Queensland fruit flies provided raspberry ketone supplements as immature adults. *PLOS ONE* 12(8): e0184086. <https://doi.org/10.1371/journal.pone.0184086>
Correction (2018): *PLOS ONE* 13(1): e0191145. <https://doi.org/10.1371/journal.pone.0191145>
- Akter, H., V. Mendez, R. Morelli, J. Pérez, and P. W. Taylor. 2017b.** Raspberry ketone supplement promotes early sexual maturation in male Queensland fruit fly, *Bactrocera tryoni* (Diptera: Tephritidae). *Pest Management Science* 73: 1764–1770. <https://doi.org/10.1002/ps.4538>
- Aluja, M., I. Jácome, and R. Macías-Ordóñez. 2001.** Effect of adult nutrition on male sexual performance in four neotropical fruit fly species of the genus *Anastrepha* (Diptera: Tephritidae). *Journal of Insect Behavior* 14: 759–775. <https://doi.org/10.1023/A:1013037400676>
- Aluja, M., M. Ordano, P. E. A. Teal, J. Sivinski, D. García-Medel, and A. Anzués-Dadda. 2009.** Larval feeding substrate and species significantly influence the effect of a juvenile hormone analog on sexual development/performance in four tropical tephritid flies. *Journal of Insect Physiology* 55: 231–242. <https://doi.org/10.1016/j.jinsphys.2008.11.013>
- Andress, E., E. Jones, M. War, and T. Shelly. 2012.** Effects of pre-release chilling on the flight ability of sterile males of the Mediterranean fruit fly (Diptera: Tephritidae). *Florida Entomologist* 95(3): 587–592. <https://doi.org/10.1653/024.095.0308>
- Andress, E., M. War, and T. Shelly. 2013.** Effect of pre-release storage on the flight ability of sterile Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* 96(4): 1615–1617. <https://doi.org/10.1653/024.096.0451>
- Andress, E., M. War, and T. Shelly. 2015.** Effect of pupal holding density on emergence rate, flight ability, and yield of sterile male Mediterranean fruit flies (Diptera: Tephritidae). *Proceedings of the Hawaiian Entomological Society* 47: 27–34. <https://pdfs.semanticscholar.org/ca96/449a3d02b3cee373aa96e6f849fc8c2f7ba8.pdf>
- Arredondo, J., L. Ruiz, E. Hernández, P. Montoya, and F. Díaz-Fleischer. 2016.** Comparison of *Anastrepha ludens* (Diptera: Tephritidae) bisexual and genetic sexing (Tapachula-7) strains: effect of hypoxia, fly density, chilling period, and food type on fly quality. *Journal of Economic Entomology* 109: 572–579. <https://doi.org/10.1093/jee/tov350>
- Augustinos, A. A., G. A. Kyritsis, N. T. Papadopoulos, A. M. M. Abd-Alla, C. Cáceres, and K. Bourtzis. 2015.** Exploitation of the medfly gut microbiota for the enhancement of sterile insect technique: use of *Enterobacter* sp. in larval diet-based probiotic applications. *PLOS ONE* 10(9): e0136459. <https://doi.org/10.1371/journal.pone.0136459>
- Bachmann, G. E., D. F. Segura, F. Devescovi, M. L. Juárez, M. J. Ruiz, M. T. Vera, J. L. Cladera, P. E. A. Teal, and P. C. Fernández. 2015.** Male sexual behavior and pheromone emission is enhanced by exposure to guava fruit volatiles in *Anastrepha fraterculus*. *PLOS ONE* 10(4): e0124250. <https://doi.org/10.1371/journal.pone.0124250>
Correction (2015): *PLOS ONE* 10(5): e0129523. <https://doi.org/10.1371/journal.pone.0129523>
- Bachmann, G. E., F. Devescovi, A. L. Nussenbaum, J. L. Cladera, P. C. Fernández, M. T. Vera, P. E. A. Teal, and D. F. Segura. 2017.** Male sexual enhancement after methoprene treatment in *Anastrepha fraterculus* (Diptera: Tephritidae): a sustained response that does not fade away after sexual maturation. *J. of Insect Physiology* 101: 7–14. <https://doi.org/10.1016/j.jinsphys.2017.06.009>

- Barclay, H. J., D. McInnis, and J. Hendrichs. 2014. Modeling the area-wide integration of male annihilation and the simultaneous release of methyl eugenol-exposed *Bactrocera* spp. sterile males. *Annals of the Entomological Society of America* 107: 97–112. <https://doi.org/10.1603/AN13010>
- Behar, A., B. Yuval, and E. Jurkevitch. 2005. Enterobacteria-mediated nitrogen fixation in natural populations of the fruit fly *Ceratitis capitata*. *Molecular Ecology* 14(9): 2637–2643. <https://doi.org/10.1111/j.1365-294X.2005.02615.x>
- Behar, A., E. Jurkevitch, and B. Yuval. 2008a. Bringing back the fruit into fruit fly-bacteria interactions. *Molecular Ecology* 17(5): 1375–1386. <https://doi.org/10.1111/j.1365-294X.2008.03674.x>
- Behar, A., B. Yuval, and E. Jurkevitch. 2008b. Gut bacterial communities in the Mediterranean fruit fly (*Ceratitis capitata*) and their impact on host longevity. *Journal of Insect Physiology* 54(9): 1377–1383. <https://doi.org/10.1016/j.jinsphys.2008.07.011>
- Behar, A., B. Yuval, and E. Jurkevitch. 2008c. Community structure of the Mediterranean fruit fly microbiota: seasonal and spatial sources of variation. *Israel Journal of Ecology and Evolution* 54(2): 181–191. <https://doi.org/10.1080/15659801.2008.10639612>
- Ben Ami, E., B. Yuval, and E. Jurkevitch. 2010. Manipulation of the microbiota of mass-reared Mediterranean fruit flies *Ceratitis capitata* (Diptera: Tephritidae) improves sterile male sexual performance. *The International Society for Microbial Ecology (ISME) Journal* 4: 28–37. <https://doi.org/10.1038/ismej.2009.82>
- Ben-Yosef, M., A. Behar, E. Jurkevitch, and B. Yuval. 2008a. Bacteria-diet interactions affect longevity in the medfly – *Ceratitis capitata*. *Journal of Applied Entomology* 132(9–10): 690–694. <https://doi.org/10.1111/j.1439-0418.2008.01330.x>
- Ben-Yosef, M., E. Jurkevitch, and B. Yuval. 2008b. Effect of bacteria on nutritional status and reproductive success of the Mediterranean fruit fly *Ceratitis capitata*. *Physiological Entomology* 33: 145–154. <https://doi.org/10.1111/j.1365-3032.2008.00617.x>
- Ben-Yosef, M., Y. Aharon, E. Jurkevitch, and B. Yuval. 2010. Give us the tools and we will do the job: symbiotic bacteria affect olive fly fitness in a diet-dependent fashion. *Proceedings of the Royal Society B. Biological Sciences* 277(1687): 1545–1552. <https://doi.org/10.1098/rspb.2009.2102>
- Ben-Yosef, M., Z. Pasternak, E. Jurkevitch, and B. Yuval. 2014. Symbiotic bacteria enable olive flies (*Bactrocera oleae*) to exploit intractable sources of nitrogen. *Journal of Evolutionary Biology* 27: 2695–2705. <https://onlinelibrary.wiley.com/doi/pdf/10.1111/jeb.12527>
- Ben-Yosef, M., Z. Pasternak, E. Jurkevitch, and B. Yuval. 2015. Symbiotic bacteria enable olive fly larvae to overcome host defences. *Royal Society Open Science* 2(7): 150170. <https://doi.org/10.1098/rsos.150170>
- Blay, S., and B. Yuval. 1997. Nutritional correlates to reproductive success of male Mediterranean fruit flies (Diptera: Tephritidae). *Animal Behaviour* 54: 59–66. <https://doi.org/10.1006/anbe.1996.0445>
- Briceño, D., W. Eberhard, and T. Shelly. 2007. Male courtship behavior in *Ceratitis capitata* (Diptera: Tephritidae) that have received aromatherapy with ginger root oil. *Florida Entomologist* 90: 175–179. [https://doi.org/10.1653/0015-4040\(2007\)90\[175:MCBICC\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[175:MCBICC]2.0.CO;2)
- Cáceres, C., G. Tsiamis, B. Yuval, E. Jurkevitch, and K. Bourtzis. 2019. Joint FAO/IAEA coordinated research project on “use of symbiotic bacteria to reduce mass-rearing costs and increase mating success in selected fruit pests in support of SIT application”. *BMC Microbiology* 19: 284. <https://doi.org/10.1186/s12866-019-1644-y>
- Cai, Z., Z. Yao, Y. Li, Z. Xi, K. Bourtzis, Z. Zhao, S. Bai, and H. Zhang. 2018. Intestinal probiotics restore the ecological fitness decline of *Bactrocera dorsalis* by irradiation. *Evolutionary Applications* 11(10): 1946–1963. <https://doi.org/10.1111/eva.12698>
- Collins, S. R., O. L. Reynolds, and P. W. Taylor. 2014. Combined effects of dietary yeast supplementation and methoprene treatment on sexual maturation of Queensland fruit fly. *Journal of Insect Physiology* 61: 51–57. <https://doi.org/10.1016/j.jinsphys.2014.01.002>
- Cunningham, R. T. 1989. Parapheromones, pp. 221–230. In A. S. Robinson and G. Hooper (eds.), *Fruit flies: their biology, natural enemies and control*. Vol. 3A. Elsevier, Amsterdam, The Netherlands.
- Damodaram, K. J. P., A. Ayyasamy, and V. Kempraj. 2016. Commensal bacteria aid mate-selection in the fruit fly, *Bactrocera dorsalis*. *Microbial Ecology* 72: 725–729. <https://doi.org/10.1007/s00248-016-0819-4>
- Deutscher, A. T., T. A. Chapman, L. A. Shuttlesworth, M. Riegler, and O. L. Reynolds. 2019. Tephritid-microbial interactions to enhance fruit fly performance in sterile insect technique programs. *BMC Microbiology* 19(Suppl. 1): 287. <https://doi.org/10.1186/s12866-019-1650-0>

- Díaz-Fleischer, F., J. Arredondo, and M. Aluja. 2009.** Enriching early adult environment affects the copulation behaviour of a tephritid fly. *Journal of Experimental Biology* 212: 2120–2127. <https://jeb.biologists.org/content/212/13/2120>
- DIR-SIT. 2020.** Directory of sterile insect technique facilities. FAO/IAEA, Vienna, Austria. [https://nucleus.iaea.org/sites/naipc/dirsit/SitePages/World-Wide%20Directory%20of%20SIT%20Facilities%20\(DIR-SIT\).aspx](https://nucleus.iaea.org/sites/naipc/dirsit/SitePages/World-Wide%20Directory%20of%20SIT%20Facilities%20(DIR-SIT).aspx)
- Epsky, N. D., and R. R. Heath. 1993.** Food availability and pheromone production by males of *Anastrepha suspensa* (Diptera: Tephritidae). *Environmental Entomology* 22: 942–947. <https://doi.org/10.1093/ee/22.5.942>
- (FAO/IAEA) Food and Agriculture Organization of the United Nations/International Atomic Energy Agency. 2017.** Guideline for packing, shipping, holding and release of sterile flies in area-wide fruit fly control programmes. J. L. Zavala-López and W. R. Enkerlin (eds.). Second Edition. FAO, Rome, Italy. 140 pp. <https://www.iaea.org/resources/manual/guideline-for-packing-shipping-holding-and-release-of-sterile-flies-in-area-wide-fruit-fly-control-programmes>
- (FAO/IAEA) Food and Agriculture Organization of the United Nations/International Atomic Energy Agency. 2018.** “The dashboard” for managers of sterile insect technique production facilities. A mass-rearing metrics and monitoring tool. P. Rendón, A. Aldana and C. Cáceres (eds.), Vienna, Austria. 18 pp. <http://www.naweb.iaea.org/nafa/ipc/public/MANUAL-MASS-REARING-METRICS-AND-MONITORING-TOOL-Ver-1.pdf>
- (FAO/IAEA) Food and Agriculture Organization of the United Nations/International Atomic Energy Agency. 2019.** Sterile insect release density calculations spreadsheet. P. Rendón, W. R. Enkerlin and C. Cáceres (eds.). Vienna, Austria. 30 pp. <http://www.naweb.iaea.org/nafa/ipc/public/RELEASE-DENSITIES-MANUAL-V.2.0.pdf>
- (FAO/IAEA/USDA) Food and Agriculture Organization of the United Nations/International Atomic Energy Agency/United States Department of Agriculture. 2019.** Product quality control for sterile mass-reared and released tephritid fruit flies. Version 7.0. IAEA, Vienna, Austria. 148 pp. <https://www.iaea.org/sites/default/files/qcv7.pdf>
- Faria, M. J., R. Pereira, T. Dellinger, and P. E. A. Teal. 2008.** Influence of methoprene and protein on survival, maturation and sexual performance of male *Ceratitis capitata* (Diptera: Tephritidae). *Journal of Applied Entomology* 132: 812–819. <https://doi.org/10.1111/j.1439-0418.2008.01360.x>
- Fezza, T. J., and T. E. Shelly. 2018.** Raspberry ketone-supplemented diet has no effect on fitness parameters or lure responsiveness in male melon flies, *Zeugodacus cucurbitae* (Diptera: Tephritidae). *Journal of Asia-Pacific Entomology* 21: 1384–1388. <https://doi.org/10.1016/j.aspen.2018.10.017>
- Flores, S., J. P. Rivera, E. Hernandez, and P. Montoya. 2011.** The effect of ginger oil on the sexual performance of *Anastrepha* males (Diptera: Tephritidae). *Florida Entomologist* 94(4): 916–922. <https://doi.org/10.1653/024.094.0428>
- Gavriel, S., Y. Gazit, and B. Yuval. 2009.** Remating by female Mediterranean fruit flies (*Ceratitis capitata*, Diptera: Tephritidae): temporal patterns and modulation by male condition. *Journal of Insect Physiology* 55: 637–642. <https://doi.org/10.1016/j.jinsphys.2009.04.002>
- Gavriel, S., Y. Gazit, and B. Yuval. 2010.** Effect of diet on survival, in the laboratory and the field, of sterile male Mediterranean fruit flies. *Entomologia Experimentalis et Applicata* 135: 96–104. <https://doi.org/10.1111/j.1570-7458.2010.00972.x>
- Gavriel, S., E. Jurkevitch, Y. Gazit, and B. Yuval. 2011.** Bacterially enriched diet improves sexual performance of sterile male Mediterranean fruit flies. *Journal of Applied Entomology* 135: 564–573. <https://doi.org/10.1111/j.1439-0418.2010.01605.x>
- Gerofotis, C. D., C. S. Ioannou, and N. T. Papadopoulos. 2013.** Aromatized to find mates: α -pinene aroma boosts the mating success of adult olive fruit flies. *PLOS ONE* 8(11): e81336. <https://doi.org/10.1371/journal.pone.0081336>
- (GFV) Good Fruit & Vegetables. 2020.** Caffeine helps fruit flies get on with gettin' it on. <https://www.goodfruitandvegetables.com.au/story/6614015/coffee-helps-q-flies-in-fertility-futility/>
- Gómez, Y., and P. E. A. Teal. 2010.** Hormonas juveniles y su aplicación en la Técnica del Insecto Estéril, pp. 357–368. In P. Montoya, J. Toledo and E. Hernández (eds.), *Moscas de la fruta: fundamentos y procedimientos para su manejo*. S y G Editores, México D.F., Mexico.
- Gómez, Y., P. E. A. Teal, and R. Pereira. 2013.** Enhancing efficacy of Mexican fruit fly SIT programmes by large-scale incorporation of methoprene into pre-release diet. *Journal of Applied Entomology* 137(Issue s1): 252–259. <https://doi.org/10.1111/j.1439-0418.2011.01695.x>

- Gomez-Simuta, Y., F. Diaz-Fleisher, J. Arredondo, E. Díaz-Santiz, and D. Pérez-Staples. 2017. Precocious Mexican fruit fly methoprene-fed males inhibit female receptivity and perform sexually as mature males. *Journal of Applied Entomology* 141: 266–273. <https://doi.org/10.1111/jen.12334>
- González-López, G. I., D. Rao, F. Díaz-Fleischer, D. Orozco-Dávila, and D. Pérez-Staples. 2016. Antipredator behavior of the new mass-reared unisexual strain of the Mexican fruit fly. *Bulletin of Entomological Research* 106(3): 314–321. <https://doi.org/10.1017/S0007485315000966>
- Hamden, H., M. M. Guerfali, S. Fadhl, M. Saidi, and C. Chevrier. 2013. Fitness improvement of mass-reared sterile males of *Ceratitis capitata* (Vienna 8 strain) (Diptera: Tephritidae) after gut enrichment with probiotics. *Journal of Economic Entomology* 106: 641–647. <https://doi.org/10.1603/EC12362>
- Haq, I., and J. Hendrichs. 2013. Pre-release feeding on hydrolysed yeast and methoprene treatment enhances male *Bactrocera cucurbitae* Coquillett (Diptera: Tephritidae) longevity. *Journal of Applied Entomology* 137(Issue s1): 99–102. <https://doi.org/10.1111/j.1439-0418.2011.01617.x>
- Haq, I. U., C. Cáceres, J. Hendrichs, P. E. A. Teal, V. Wornoayporn, C. Stauffer, and A. S. Robinson. 2010a. Effects of the juvenile hormone analogue methoprene and dietary protein on male melon fly *Bactrocera cucurbitae* (Diptera:Tephritidae) mating success. *Journal of Insect Physiology* 56: 1503–1509. <https://doi.org/10.1016/j.jinsphys.2010.04.018>
- Haq, I. U., C. Cáceres, J. Hendrichs, P. E. A. Teal, C. Stauffer, and A. S. Robinson. 2010b. Methoprene modulates the effect of diet on male melon fly, *Bactrocera cucurbitae*, performance at mating aggregations. *Entomologia Experimentalis et Applicata* 136: 21–30. <https://doi.org/10.1111/j.1570-7458.2010.00999.x>
- Haq, I. U., L. Mayr, P. E. A. Teal, J. Hendrichs, A. S. Robinson, C. Stauffer, and R. Hood-Nowotny. 2010c. Total body nitrogen and total body carbon as indicators of body protein and body lipids in the melon fly *Bactrocera cucurbitae*: effects of methoprene, a juvenile hormone analogue, and of diet supplementation with hydrolyzed yeast. *Journal of Insect Physiology* 56: 1807–1815. <https://doi.org/10.1016/j.jinsphys.2010.07.011>
- Haq, I., C. Cáceres, P. Liedo, D. Soriano, A. Jessup, J. Hendrichs, P. E. A. Teal, and A. S. Robinson. 2013. Effect of methoprene application, adult food and feeding duration on male melon fly starvation survival. *Journal of Applied Entomology* 137(Issue s1): 61–68. <https://doi.org/10.1111/j.1439-0418.2010.01575.x>
- Haq, I. U., M. J. B. Vreysen, P. E. A. Teal, and J. Hendrichs. 2014a. Methoprene application and diet protein supplementation to male melon fly, *Bactrocera cucurbitae*, modifies female remating behavior. *Insect Science* 21: 637–646. <https://doi.org/10.1111/1744-7917.12073>
- Haq, I., M. J. B. Vreysen, C. Cáceres, T. E. Shelly, and J. Hendrichs. 2014b. Methyl eugenol aromatherapy enhances the mating competitiveness of male *Bactrocera carambolae* Drew & Hancock (Diptera: Tephritidae). *Journal of Insect Physiology* 68: 1–6. <https://doi.org/10.1016/j.jinsphys.2014.06.014>
- Haq, I. U., M. J. B. Vreysen, C. Cáceres, T. E. Shelly, and J. Hendrichs. 2015. Optimizing methyl-eugenol aromatherapy to maximize posttreatment effects to enhance mating competitiveness of male *Bactrocera carambolae* (Diptera: Tephritidae). *Insect Science* 22: 661–669. <https://doi.org/10.1111/1744-7917.12148>
- Haq, I. U., C. Cáceres, J. S. Meza, J. Hendrichs, and M. J. B. Vreysen. 2018. Different methods of methyl eugenol application enhance the mating success of male oriental fruit fly (Diptera: Tephritidae). *Scientific Reports* 8: 6033. <https://doi.org/10.1038/s41598-018-24518-5>
- Hee, A. K. W., and K. H. Tan. 1998. Attraction of female and male *Bactrocera papayae* to conspecific males fed with methyl eugenol and attraction of females to male sex pheromone components. *Journal of Chemical Ecology* 24: 753–764. <https://doi.org/10.1023/A:1022302605357>
- Hendrichs, J., and M. A. Hendrichs. 1990. Mediterranean fruit fly (Diptera: Tephritidae) in nature: location and diel pattern of feeding and other activities on fruiting and nonfruiting hosts and nonhosts. *Annals of the Entomological Society of America* 83: 632–641. <https://doi.org/10.1093/aesa/83.3.632>
- Hendrichs, J., and R. J. Prokopy. 1994. Food foraging behavior of frugivorous fruit flies, pp. 37–55. In C. O. Calkins, W. Klassen and P. Liedo (eds.), *Fruit flies and the sterile insect technique*. CRC Press, Boca Raton, FL, USA.
- Hendrichs, J., B. I. Katsoyannos, D. R. Papaj, and R. J. Prokopy. 1991. Sex differences in movement between natural feeding and mating sites and tradeoffs between food consumption, mating success and predator evasion in Mediterranean fruit flies (Diptera: Tephritidae). *Oecologia* 86: 223–231. <https://doi.org/10.1007/BF00317534>

- Hendrichs, M. A., and J. Hendrichs. 1998.** Perfumed to be killed: interception of Mediterranean fruit fly (Diptera: Tephritidae) sexual signaling by predatory foraging wasps (Hymenoptera: Vespidae). *Annals of the Entomological Society of America* 91: 228–234. <https://doi.org/10.1093/aesa/91.2.228>
- Inskeep, J. R., T. E. Shelly, R. I. Vargas, and H. Spafford. 2019.** Zingerone feeding affects mate choice but not fecundity or fertility in the melon fly, *Zeugodacus cucurbitae* (Diptera: Tephritidae). *Florida Entomologist* 102: 161–167. <https://doi.org/10.1653/024.102.0126>
- Ji, Q. E., J. H. Chen, D. O. McInnis, and Q. L. Guo. 2013.** The effect of methyl eugenol exposure on subsequent mating performance of sterile males of *Bactrocera dorsalis*. *Journal of Applied Entomology* 137(Issue s1): 238–243. <https://doi.org/10.1111/j.1439-0418.2011.01686.x>
- Juan-Blasco, M., V. San Andrés, M. A. Martínez-Utrillas, R. Argilés, I. Pla, A. Urbaneja, and B. Sabater-Muñoz. 2013.** Alternatives to ginger root oil aromatherapy for improved mating performance of sterile *Ceratitis capitata* (Diptera: Tephritidae) males. *Journal of Applied Entomology* 137(Issue s1): 244–251. <https://doi.org/10.1111/j.1439-0418.2011.01688.x>
- Juárez, M. L., L. E. Pimper, G. E. Bachmann, C. A. Conte, M. J. Ruiz, L. Goane, P. Medina Pereyra, F. Castro, J. Salgueiro, J. L. Cladera, P. C. Fernández, K. Bourtzis, S. B. Lanzavecchia, M. T. Vera, and D. F. Segura. 2019.** Gut bacterial diversity and physiological traits of *Anastrepha fraterculus* Brazilian-1 morphotype males are affected by antibiotic treatment. *BMC Microbiology* 19: 283. <https://doi.org/10.1186/s12866-019-1645-x>
- Kaspi, R., and B. Yuval. 2000.** Post-teneral protein feeding improves sexual competitiveness but reduces longevity of mass-reared sterile male Mediterranean fruit flies (Diptera: Tephritidae). *Annals of the Entomological Society of America* 93: 949–955. [https://doi.org/10.1603/0013-8746\(2000\)093\[0949:PTPFIJ\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2000)093[0949:PTPFIJ]2.0.CO;2)
- Katsoyannos, B. I., N. A. Kouloussis, and N. T. Papadopoulos. 1997.** Response of *Ceratitis capitata* to citrus chemicals under semi-natural conditions. *Entomologia Experimentalis et Applicata* 82: 181–188. <https://doi.org/10.1046/j.1570-7458.1997.00129.x>
- Katsoyannos, B. I., N. T. Papadopoulos, N. K. Kouloussis, and J. Hendrichs. 2004.** Effect of citrus peel substances on male Mediterranean fruit fly behaviour, pp. 13–17. In B. N. Barnes (ed.), *Proceedings of the 6th International Symposium on Fruit Flies of Economic Importance*, 6–10 May 2002, Stellenbosch, South Africa. Isteg Scientific Publications, Irene, South Africa. https://nucleus.iaea.org/sites/naipc/twd/Documents/6thISFFEI_Proceedings/KATSOYANNOS.pdf
- Khan, M. A. M., L. A. Shuttleworth, T. Osborne, D. Collins, G. M. Gurr, and O. L. Reynolds. 2019.** Raspberry ketone accelerates sexual maturation and improves mating performance of sterile male Queensland fruit fly, *Bactrocera tryoni* (Froggatt). *Pest Management Science* 75(7): 1942–1950. <https://doi.org/10.1002/ps.5307>
- Khoo, C. C. H., and K. H. Tan. 2000.** Attraction of both sexes of melon fly, *Bactrocera cucurbitae* to conspecific males - a comparison after pharmacophagy of cue-lure and a new attractant - zingerone. *Entomologia Experimentalis et Applicata* 97: 317–320. <https://doi.org/10.1046/j.1570-7458.2000.00745.x>
- Kokkari, A. I., O. D. Pliakou, G. D. Floros, N. A. Kouloussis, and D. S. Koveos. 2017.** Effect of fruit volatiles and light intensity on the reproduction of *Bactrocera (Dacus) oleae*. *Journal of Applied Entomology* 141: 841–847. <https://doi.org/10.1111/jen.12389>
- Kouloussis, N. A., B. I. Katsoyannos, N. T. Papadopoulos, C. S. Ioannou, and I. V. Iliadis. 2013.** Enhanced mating competitiveness of *Ceratitis capitata* males following exposure to citrus compounds. *Journal of Applied Entomology* 137(Issue s1): 30–38. <https://doi.org/10.1111/j.1439-0418.2010.01535.x>
- Kouloussis, N. A., C. D. Geroftis, C. S. Ioannou, I. V. Iliadis, N. T. Papadopoulos, and D. S. Koveos. 2017.** Towards improving sterile insect technique: exposure to orange oil compounds increases sexual signalling and longevity in *Ceratitis capitata* males of the Vienna 8 GSS. *PLOS ONE* 12(11): e0188092. <https://doi.org/10.1371/journal.pone.0188092>
- Kumaran, N., S. Balagawi, M. K. Schutze, and A. R. Clarke. 2013.** Evolution of lure response in tephritid fruit flies: phytochemicals as drivers of sexual selection. *Animal Behaviour* 85: 781–789. <https://doi.org/10.1016/j.anbehav.2013.01.024>
- Kumaran, N., P. J. Prentis, K. P. Mangalam, M. K. Schutze, and A. R. Clarke. 2014a.** Sexual selection in true fruit flies (Diptera: Tephritidae): transcriptome and experimental evidences for phytochemicals increasing male competitive ability. *Molecular Ecology* 23(18): 4645–4657. <https://doi.org/10.1111/mec.12880>

- Kumaran, N., R. A. Hayes, and A. R. Clarke. 2014b. Cuelure but not zingerone make the sex pheromone of male *Bactrocera tryoni* (Tephritidae: Diptera) more attractive to females. *Journal of Insect Physiology* 68: 36–43. <https://doi.org/10.1016/j.jinsphys.2014.06.015>
- Kyritsis, G. A., A. A. Augustinos, S. Ntougias, N. T. Papadopoulos, K. Bourtzis, and C. Cáceres. 2019. *Enterobacter* sp. AA26 gut symbiont as a protein source for Mediterranean fruit fly mass-rearing and sterile insect technique applications. *BMC Microbiology* 19: 288. <https://doi.org/10.1186/s12866-019-1651-z>
- Levy, K., T. E. Shelly, and B. Yuval. 2005. Effects of the olfactory environment and nutrition on the ability of male Mediterranean fruit flies to endure starvation. *Journal of Economic Entomology* 98: 61–65. <https://doi.org/10.1093/jee/98.1.61>
- Liedo, P., E. De Leon, M. I. Barrios, J. F. Valle-Mora, and G. Ibarra. 2002. Effect of age on the mating propensity of the Mediterranean fruit fly (Diptera Tephritidae). *Florida Entomologist* 85: 94–101. [https://doi.org/10.1653/0015-4040\(2002\)085\[0094:EOAOTM\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2002)085[0094:EOAOTM]2.0.CO;2)
- Liedo, P., D. Orozco, L. Cruz-López, J. L. Quintero, C. Becerra-Pérez, M. D. R. Hernández, A. Oropeza, and J. Toledo. 2013. Effect of post-teneral diets on the performance of sterile *Anastrepha ludens* and *Anastrepha obliqua* fruit flies. *Journal of Applied Entomology* 137(Issue s1): 49–60. <https://doi.org/10.1111/j.1439-0418.2010.01568.x>
- Liendo, M. C., F. Devescovi, G. E. Bachmann, M. E. Utgés, S. Abraham, M. T. Vera, S. B. Lanzavecchia, J. P. Bouvet, P. Gómez-Cendra, J. Hendrichs, P. E. A. Teal, J. L. Cladera, and D. F. Segura. 2013. Precocious sexual signalling and mating in *Anastrepha fraterculus* (Diptera: Tephritidae) sterile males achieved through juvenile hormone treatment and protein supplements. *Bulletin of Entomological Research* 103: 1–13. <https://doi.org/10.1017/S0007485312000442>
- Maor, M., B. Kamensky, S. Shloush, and B. Yuval. 2004. Effects of post-teneral diet on foraging success of sterile male Mediterranean fruit flies. *Entomologia Experimentalis et Applicata* 110: 225–230. <https://doi.org/10.1111/j.0013-8703.2004.00139.x>
- McInnis, D., R. Kurashima, T. Shelly, J. Komatsu, J. Edu, and E. Pahio. 2011. Prerelease exposure to methyl eugenol increases the mating competitiveness of sterile males of the oriental fruit fly (Diptera: Tephritidae) in a Hawaiian orchard. *Journal of Economic Entomology* 104: 1969–1978. <https://doi.org/10.1603/EC11134>
- McInnis, D. O., B. J. Paranhos, and T. E. Shelly. 2013. Survival of sterile male Mediterranean fruit flies in large field cages after release at different ages. *Journal of Applied Entomology* 137(Issue s1): 43–48. <https://doi.org/10.1111/j.1439-0418.2010.01567.x>
- Morató, S., T. Shelly, J. Rull, and M. Aluja. 2015. Sexual competitiveness of *Anastrepha ludens* (Diptera: Tephritidae) males exposed to *Citrus aurantium* and *Citrus paradisi* essential oils. *Journal of Economic Entomology* 108: 621–628. <https://doi.org/10.1093/jee/tou054>
- Morelli, R., B. J. Paranhos, A. M. Coelho, R. Castro, L. Garziera, F. Lopes, and J. M. S. Bento. 2013. Exposure of sterile Mediterranean fruit fly (Diptera: Tephritidae) males to ginger root oil reduces female remating. *Journal of Applied Entomology* 137(Issue s1): 75–82. <https://doi.org/10.1111/j.1439-0418.2010.01584.x>
- Morimoto, J., B. Nguyen, S. T. Tabrizi, I. Lundbäck, P. W. Taylor, F. Ponton, and T. A. Chapman. 2019. Commensal microbiota modulates larval foraging behaviour, development rate and pupal production in *Bactrocera tryoni*. *BMC Microbiology* 19: 286. <https://doi.org/10.1186/s12866-019-1648-7>
- Ndzana, A. R. F. X., T. Turlings, N. Woin, and S. Quilici. 2016. Factors influencing the mating success of *Bactrocera zonata* (Saunders) (Diptera: Tephritidae) males in an SIT programme. In *Third International Symposium on Tephritid Workers of Europe, Africa and the Middle East*. 11–14 April 2016, Stellenbosch, South Africa. <https://agritrop.cirad.fr/583108/> <http://teamfly2016.co.za/abstracts/>
- Niyazi, N., C. R. Lauzon, and T. E. Shelly. 2004. Effect of probiotic adult diets on fitness components of sterile male Mediterranean fruit flies (Diptera: Tephritidae) under laboratory and field cage conditions. *Journal of Economic Entomology* 97: 1570–1580. <https://doi.org/10.1603/0022-0493-97.5.1570>
- Noman, M. S., L. Liu, Z. Bai, and Z. Li. 2020. Tephritidae bacterial symbionts: potentials for pest management. *Bulletin of Entomological Research* 110: 1–14. <https://doi.org/10.1017/S0007485319000403>
- Obra, G. B., and S. S. Resilva. 2013. Influence of adult diet and exposure to methyl eugenol in the mating performance of *Bactrocera philippinensis*. *Journal of Applied Entomology* 137(Issue s1): 210–216. <https://doi.org/10.1111/j.1439-0418.2011.01678.x>

- Orankanok, W., S. Chinvinijkul, A. Sawatwangkhong, S. Pinkaew, and S. Orankanok. 2013. Methyl eugenol and pre-release diet improve mating performance of young *Bactrocera dorsalis* and *Bactrocera correcta* males. *Journal of Applied Entomology* 137(Issue s1): 200–209. <https://doi.org/10.1111/j.1439-0418.2011.01677.x>
- Papadopoulos, N. T., B. I. Katsoyannos, N. A. Kouloussis, and J. Hendrichs. 2001. Effect of orange peel substances on mating competitiveness of male *Ceratitis capitata*. *Entomologia Experimentalis et Applicata* 99: 253–261. <https://doi.org/10.1046/j.1570-7458.2001.00824.x>
- Papadopoulos, N. T., T. E. Shelly, N. Niyazi, and E. Jang. 2006. Olfactory and behavioral mechanisms underlying enhanced mating competitiveness following exposure to ginger root oil and orange oil in males of the Mediterranean fruit fly, *Ceratitis capitata* (Diptera: Tephritidae). *Journal of Insect Behavior* 19: 403. <https://doi.org/10.1007/s10905-006-9031-6>
- Paranhos, B. J., D. McInnis, K. Uramoto, I. Damasceno, N. Gonçalves, R. M. Alves, M. L. Costa, J. Walder, A. Malavasi, and A. Nascimento. 2008. Sterile medfly males of the TSL Vienna 8 genetic sexing strain display improved mating performance with ginger root oil, pp. 313–318. In R. L. Sugayama, R. A. Zucchi, S. M. Ovruski and J. Sivinski (eds.), *Fruit flies of economic importance: from basic to applied knowledge*. Sociedade Brasileira para o Progresso da Ciência (SBPC). Salvador, Brazil.
- Paranhos, B. J., N. T. Papadopoulos, D. McInnis, C. Gava, F. S. C. Lopes, R. Morelli, and A. Malavasi. 2010. Field dispersal and survival of sterile medfly males aromatically treated with ginger root oil. *Environmental Entomology* 39: 570–575. <https://doi.org/10.1603/EN08309>
- Paranhos, B. J., D. McInnis, R. Morelli, R. M. Castro, L. Garziera, L. G. Paranhos, K. Costa, C. Gava, M. L. Z. Costa, and J. M. M. Walder. 2013. Optimum dose of ginger root oil to treat sterile Mediterranean fruit fly males (Diptera: Tephritidae). *Journal of Applied Entomology* 137(Issue s1): 83–90. <https://doi.org/10.1111/j.1439-0418.2010.01595.x>
- Pereira, R., J. Sivinski, and P. E. A. Teal. 2009. Influence of methoprene and dietary protein on male *Anastrepha suspensa* (Diptera: Tephritidae) mating aggregations. *Journal of Insect Physiology* 55: 328–335. <https://doi.org/10.1016/j.jinsphys.2008.12.010>
- Pereira, R., J. Sivinski, and P. E. A. Teal. 2010a. Influence of a juvenile hormone analog and dietary protein on male *Anastrepha suspensa* (Diptera: Tephritidae) sexual success. *Journal of Economic Entomology* 103: 40–46. <https://doi.org/10.1603/EC08324>
- Pereira, R., J. Sivinski, P. Teal, and J. Brockmann. 2010b. Enhancing male sexual success in a lekking fly (*Anastrepha suspensa* Diptera: Tephritidae) through a juvenile hormone analog has no effect on adult mortality. *Journal of Insect Physiology* 56: 1552–1557. <https://doi.org/10.1016/j.jinsphys.2010.05.008>
- Pereira, R., J. Sivinski, J. P. Shapiro, and P. E. A. Teal. 2011. Influence of methoprene and dietary protein on male *Anastrepha suspensa* (Diptera: Tephritidae) lipid and protein content. *Florida Entomologist* 94: 137–144. <https://doi.org/10.1653/024.094.0203>
- Pereira, R., B. Yuval, P. Liedo, P. E. A. Teal, T. E. Shelly, D. O. McInnis, and J. Hendrichs. 2013a. Improving sterile male performance in support of programmes integrating the sterile insect technique against fruit flies. *Journal of Applied Entomology* 137(Issue s1): 178–190. <https://doi.org/10.1111/j.1439-0418.2011.01664.x>
- Pereira, R., P. E. A. Teal, H. Conway, J. Worley, and J. Sivinski. 2013b. Influence of methoprene and dietary protein on maturation and sexual performance of sterile *Anastrepha ludens* (Diptera: Tephritidae). *Journal of Applied Entomology* 137(Issue s1): 191–199. <https://doi.org/10.1111/j.1439-0418.2011.01668.x>
- Pérez-Staples, D., V. Prabhu, and P. W. Taylor. 2007. Post-teneral protein feeding enhances sexual performance of Queensland fruit flies. *Physiological Entomology* 32: 225–232. <https://doi.org/10.1111/j.1365-3032.2007.00568.x>
- Pérez-Staples, D., A. M. T. Harmer, S. R. Collins, and P. W. Taylor. 2008a. Potential for pre-release diet supplements to increase the sexual performance and longevity of male Queensland fruit flies. *Agricultural and Forest Entomology* 10: 255–262. <https://doi.org/10.1111/j.1461-9563.2008.00385.x>
- Pérez-Staples, D., M. Aluja, R. Macías-Ordóñez, and J. Sivinski. 2008b. Reproductive trade-offs from mating with a successful male: the case of the tephritid fly *Anastrepha obliqua*. *Behavioral Ecology and Sociobiology* 62: 1333–1340. <https://doi.org/10.1007/s00265-008-0561-0>
- Pérez-Staples, D., C. W. Weldon, C. Smallridge, and P. W. Taylor. 2009. Pre-release feeding on yeast hydrolysate enhances sexual competitiveness of sterile male Queensland fruit flies in field cages. *Entomologia Experimentalis et Applicata* 131: 159–166. <https://doi.org/10.1111/j.1570-7458.2009.00841.x>

- Pérez-Staples, D., C. W. Weldon, and P. W. Taylor. 2011.** Sex differences in development response to yeast hydrolysate supplements in adult Queensland fruit fly. *Entomologia Experimentalis et Applicata* 141: 103–113. <https://doi.org/10.1111/j.1570-7458.2011.01173.x>
- Prabhu, V., D. Perez-Staples, and P. W. Taylor. 2008.** Protein: carbohydrate ratios promoting sexual activity and longevity of male Queensland fruit flies. *Journal of Applied Entomology* 132: 575–582. <https://doi.org/10.1111/j.1439-0418.2007.01265.x>
- Quilici, S., P. F. Duyck, and A. Franck. 2004.** Preliminary experiments on the influence of exposure to methyl eugenol on mating success of males in the peach fruit fly, *Bactrocera zonata*. In 1st FAO/IAEA Research Coordination Meeting on Improving Sterile Male Performance in Fruit Fly SIT. 25–29 October 2004. Antigua, Guatemala. <http://www-naweb.iaea.org/nafa/ipc/crp/ipc-male-performance-in-sit.html>
- Quilici, S., C. Schmitt, J. Vidal, A. Franck, and J. P. Deguine. 2013.** Adult diet and exposure to semiochemicals influence male mating success in *Ceratitis rosa* (Diptera: Tephritidae). *Journal of Applied Entomology* 137(Issue s1): 142–153. <https://doi.org/10.1111/j.1439-0418.2011.01650.x>
- Raghu, S., and A. R. Clarke. 2003.** Spatial and temporal partitioning of behaviour by adult dacines: direct evidence for methyl eugenol as a mate rendezvous cue for *Bactrocera cacuminata*. *Physiological Entomology* 28: 175–184. <https://doi.org/10.1046/j.1365-3032.2003.00328.x>
- Rao, D., S. Aguilar-Argüello, P. Montoya, and F. Díaz-Fleischer. 2013.** The effect of irradiation and mass-rearing on the anti-predator behaviour of the Mexican fruit fly *Anastrepha ludens* (Diptera: Tephritidae). *Bulletin of Entomological Research* 104: 176–181. <https://doi.org/10.1017/S0007485313000643>
- Raza, M. F., Z. Yao, S. Bai, Z. Cai, and H. Zhang. 2020.** Tephritidae fruit fly gut microbiome diversity, function and potential for applications. *Bulletin of Entomological Research*. FirstView article. 15 pp. <https://doi.org/10.1017/S0007485319000853>
- Reyes-Hernández, M., R. Thimmappa, S. Abraham, K. J. P. Damodaram, and D. Pérez-Staples. 2019.** Methyl eugenol effects on *Bactrocera dorsalis* male total body protein, reproductive organs and ejaculate. *Journal of Applied Entomology* 143: 177–186. <https://doi.org/10.1111/jen.12576>
- Reynolds, O. L., B. A. Orchard, S. R. Collins, and P. W. Taylor. 2014.** Yeast hydrolysate supplementation increases field abundance and persistence of sexually mature sterile Queensland fruit fly, *Bactrocera tryoni* (Froggatt). *Bulletin of Entomological Research* 104: 251–261. <https://doi.org/10.1017/S0007485313000758>
- Rull, J., R. Lasa, C. Rodriguez, R. Ortega, O. E. Velazquez, and M. Aluja. 2015.** Artificial selection, pre-release diet, and gut symbiont inoculation effects on sterile male longevity for area-wide fruit-fly management. *Entomologia Experimentalis et Applicata* 157: 325–333. <https://doi.org/10.1111/eea.12371>
- Sacchetti, P., B. Ghiardi, A. Granchietti, F. M. Stefanini, and A. Belcari. 2014.** Development of probiotic diets for the olive fly: evaluation of their effects on fly longevity and fecundity. *Annals of Applied Biology* 164: 138–150. <https://doi.org/10.1111/aab.12088>
- San Andrés, V., J. Pérez-Panadés, E. A. Carbonell, P. Castañera, and A. Urbaneja. 2009.** Effects of post-teneral nutrition and ginger root oil exposure on longevity and mortality in bait treatments of sterile male *Ceratitis capitata*. *Entomologia Experimentalis et Applicata* 132: 256–263. <https://doi.org/10.1111/j.1570-7458.2009.00893.x>
- Segura, D. F., C. Cáceres, M. T. Vera, V. Wornoyaporn, A. Islam, P. E. A. Teal, J. L. Cladera, J. Hendrichs, and A. S. Robinson. 2009.** Enhancing mating performance after juvenile hormone treatment in *Anastrepha fraterculus*: a differential response in males and females acts as a physiological sexing system. *Entomologia Experimentalis et Applicata* 131: 75–84. <https://doi.org/10.1111/j.1570-7458.2009.00830.x>
- Segura, D. F., M. E. Utgés, M. C. Liendo, M. F. Rodríguez, F. Devescovi, M. T. Vera, P. E. A. Teal, and J. L. Cladera. 2013.** Methoprene treatment reduces the pre-copulatory period in *Anastrepha fraterculus* (Diptera: Tephritidae) sterile males. *Journal of Applied Entomology* 137(Issue s1): 19–29. <https://doi.org/10.1111/j.1439-0418.2010.01534.x>
- Segura, D. F., S. A. Belliard, M. T. Vera, G. E. Bachmann, M. J. Ruiz, F. Jofre-Barud, P. C. Fernández, M. L. López, and T. E. Shelly. 2018.** Plant chemicals and the sexual behavior of male tephritid fruit flies. *Annals of the Entomological Society of America* 111: 239–264. <https://doi.org/10.1093/aesa/say024>
- Shelly, T. E. 2000.** Effects of raspberry ketone on the mating success of male melon flies (Diptera: Tephritidae). *Proceedings of the Hawaiian Entomological Society* 34: 143–147. <http://hdl.handle.net/10125/8368>

- Shelly, T. E. 2001.** Exposure to α -copaene and α -copaene-containing oils enhances mating success of male Mediterranean fruit flies (Diptera: Tephritidae). *Annals of the Entomological Society of America* 94: 497–502. [https://doi.org/10.1603/0013-8746\(2001\)094\[0497:ETCACC\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2001)094[0497:ETCACC]2.0.CO;2)
- Shelly, T. E. 2008.** Aromatherapy and medfly SIT, pp. 59–69. *In* R. L. Sugayama, R. A. Zucchi, S. M. Ovruski and J. Sivinski (eds.), *Fruit flies of economic importance: from basic to applied knowledge*. Sociedade Brasileira para o Progresso da Ciência (SBPC). Salvador, Brazil.
- Shelly, T. E. 2009.** Exposure to grapefruits and grapefruit oil increases male mating success in the Mediterranean fruit fly (Diptera: Tephritidae). *Proceedings of the Hawaiian Entomological Society* 41: 31–36. <https://core.ac.uk/download/pdf/5102644.pdf>
- Shelly, T. E. 2017a.** Yeast hydrolysate deprivation and the mating success of male melon flies (Diptera: Tephritidae). *Florida Entomologist* 100: 772–776. <https://doi.org/10.1653/024.100.0401>
- Shelly, T. E. 2017b.** Zingerone and the mating success and field attraction of male melon flies (Diptera: Tephritidae). *Journal of Asia-Pacific Entomology* 20: 175–178. <https://doi.org/10.1016/j.aspen.2016.12.013>
- Shelly, T. E. 2019.** Male lures and male mating competitiveness in a genetic sexing strain of the melon fly (Diptera: Tephritidae). *Journal of Asia-Pacific Entomology* 22: 393–397. <https://doi.org/10.1016/j.aspen.2019.01.014>
- Shelly, T. E., and J. Edu. 2007.** Exposure to the ripe fruit of tropical almond enhances the mating success of male *Bactrocera dorsalis* (Diptera: Tephritidae). *Florida Entomologist* 90: 757–758. [https://doi.org/10.1653/0015-4040\(2007\)90\[757:ETTRFO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[757:ETTRFO]2.0.CO;2)
- Shelly, T. E., and J. Edu. 2008a.** Effect of nitrogen-containing dietary supplements on the mating success of sterile males of the Mediterranean fruit fly (Diptera: Tephritidae). *Florida Entomologist* 91: 394–399. [https://doi.org/10.1653/0015-4040\(2008\)91\[394:EONDSO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2008)91[394:EONDSO]2.0.CO;2)
- Shelly, T. E., and J. Edu. 2008b.** Do methyl eugenol-fed males of the oriental fruit fly (Diptera: Tephritidae) induce female re-mating? *Florida Entomologist* 91: 388–393. <https://journals.flvc.org/flaent/article/view/75824/73482>
- Shelly, T. E., and N. D. Epsky. 2015.** Exposure to tea tree oil enhances the mating success of male Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* 98: 1127–1133. <https://doi.org/10.1653/024.098.0417>
- Shelly, T. E., and S. S. Kennelly. 2003.** Starvation and the mating success of wild male Mediterranean fruit flies (Diptera: Tephritidae). *Journal of Insect Behaviour* 16: 171–179. <https://doi.org/10.1023/A:1023926717088>
- Shelly, T. E., and D. O. McInnis. 2003.** Influence of adult diet on the mating success and survival of male Mediterranean fruit flies (Diptera: Tephritidae) from two mass-rearing strains on field-caged host trees. *Florida Entomologist* 86: 340–344. [https://doi.org/10.1653/0015-4040\(2003\)086\[0340:IOADOT\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2003)086[0340:IOADOT]2.0.CO;2)
- Shelly, T. E., and J. Nishimoto. 2017.** Does female mate choice confer direct fitness benefits? Results from a tephritid fruit fly. *Annals of the Entomological Society of America* 110: 204–211. <https://doi.org/10.1093/aesa/saw073>
- Shelly, T., S. Resilva, M. Reyes, and H. Bignayan. 1996.** Methyl eugenol and mating competitiveness of irradiated male *Bactrocera philippinensis* (Diptera: Tephritidae). *Florida Entomologist* 79: 481–488. <https://journals.flvc.org/flaent/article/view/59246/56925>
- Shelly, T. E., P. Rendon, E. Hernandez, S. Salgado, D. McInnis, E. Villalobos, and P. Liedo. 2003.** Effects of diet, ginger root oil, and elevation on the mating competitiveness of male Mediterranean fruit flies (Diptera: Tephritidae) from a mass-reared, genetic sexing strain in Guatemala. *Journal of Economic Entomology* 96: 1132–1141. <https://doi.org/10.1093/jee/96.4.1132>
- Shelly, T., C. Dang, and S. Kennelly. 2004.** Exposure to orange (*Citrus sinensis* L.) trees, fruit, and oil enhances mating success of male Mediterranean fruit flies (*Ceratitis capitata* [Wiedemann]). *Journal of Insect Behavior* 17: 303–315. <https://doi.org/10.1023/B:JOIR.0000031532.29287.95>
- Shelly, T. E., J. Edu, and E. Pahio. 2005.** Influence of diet and methyl eugenol on the mating success of males of the oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *Florida Entomologist* 88: 307–313. [https://doi.org/10.1653/0015-4040\(2005\)088\[0307:IODAME\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2005)088[0307:IODAME]2.0.CO;2)
- Shelly, T. E., J. Edu, and E. Pahio. 2006a.** Dietary protein and mating competitiveness of sterile males of the Mediterranean fruit fly (Diptera: Tephritidae): measurements of induced egg sterility in large field enclosures. *Florida Entomologist* 89: 277–278. [https://doi.org/10.1653/0015-4040\(2006\)89\[277:DPAMCO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2006)89[277:DPAMCO]2.0.CO;2)

- Shelly, T. E., J. Edu, and E. Pahio. 2006b. Application of orange oil to prerelease holding boxes increases the mating success of sterile males of the Mediterranean fruit fly in field cage trials (Diptera: Tephritidae). *Proceedings of the Hawaiian Entomological Society* 38: 73–79. <http://hdl.handle.net/10125/199>
- Shelly, T. E., J. Edu, and E. Pahio. 2007a. Exposure to ginger root oil decreases capture of male Mediterranean fruit flies (Diptera: Tephritidae) in trimedlure-baited traps. *Proceedings of the Hawaiian Entomological Society* 39: 27–32. <https://pdfs.semanticscholar.org/2e37/60327995e17c83f061a72b79a7593c80b0ed.pdf>
- Shelly, T. E., D. O. McInnis, C. Rodd, J. Edu, and E. Pahio. 2007b. Sterile insect technique and Mediterranean fruit fly (Diptera: Tephritidae): assessing the utility of aromatherapy in a Hawaiian coffee field. *Journal of Economic Entomology* 100: 273–282. <https://doi.org/10.1093/jee/100.2.273>
- Shelly, T., J. Edu, E. Smith, K. Hoffman, M. War, R. Santos, A. Favela, R. Garagliano, B. Ibewiro, and D. McInnis. 2007c. Aromatherapy on a large scale: exposing entire holding rooms to ginger root oil increases the mating competitiveness of sterile males of the Mediterranean fruit fly in field cage trials. *Entomologia Experimentalis et Applicata* 123: 193–201. <https://doi.org/10.1111/j.1570-7458.2007.00538.x>
- Shelly, T. E., M. War, and A. Favela. 2008a. Exposing entire adult holding rooms containing sterile male Mediterranean fruit flies to orange oil increases the mating success of those males in field-cage trials. *Florida Entomologist* 91: 686–689. <https://www.jstor.org/stable/25434890>
- Shelly, T. E., J. Edu, E. Pahio, S. L. Wee, and R. Nishida. 2008b. Re-examining the relationship between sexual maturation and age of response to methyl eugenol in males of the oriental fruit fly. *Entomologia Experimentalis et Applicata* 128: 380–388. <https://doi.org/10.1111/j.1570-7458.2008.00710.x>
- Shelly, T. E., A. N. Cowan, J. Edu, and E. Pahio. 2008c. Mating success of male Mediterranean fruit flies following exposure to two sources of α -copaene, manuka oil and mango. *Florida Entomologist* 91: 9–15. <https://www.jstor.org/stable/20065920>
- Shelly, T. E., J. I. Nishimoto, and J. Edu. 2009. Influence of methoprene on the age-related mating propensity of males of the oriental fruit fly and the Mediterranean fruit fly (Diptera: Tephritidae). *Florida Entomologist* 92: 193–198. <https://doi.org/10.1653/024.092.0201>
- Shelly, T. E., J. Edu, and D. McInnis. 2010a. Pre-release consumption of methyl eugenol increases the mating competitiveness of sterile males of the oriental fruit fly, *Bactrocera dorsalis*, in large field enclosures. *Journal of Insect Science* 10(8): 1–16. <https://doi.org/10.1673/031.010.0801>
- Shelly, T. E., P. Rendón, F. Moscoso, and R. Menendez. 2010b. Testing the efficacy of aromatherapy at the world's largest eclosion facility for sterile males of the Mediterranean fruit fly (Diptera: Tephritidae). *Proceedings of the Hawaiian Entomological Society* 42: 33–40. <http://hdl.handle.net/10125/19915>
- Shelly, T. E., J. Edu, and J. Nishimoto. 2013. Chilling and flight ability and mating competitiveness of sterile males of the Mediterranean fruit fly. *Journal of Applied Entomology* 137(Issue s1): 11–18. <https://doi.org/10.1111/j.1439-0418.2010.01532.x>
- Silva, N., L. Dantas, R. Calisto, M. J. Faria, and R. Pereira. 2013. Improving an adult holding system for Mediterranean fruit fly, *Ceratitidis capitata*, to enhance sterile male performance. *Journal of Applied Entomology* 137(Issue s1): 230–237. <https://doi.org/10.1111/j.1439-0418.2010.01532.x>
- Sookar, P., I. Haq, A. Jessup, D. McInnis, G. Franz, V. Wornoyaporn, and S. Permalloo. 2013. Mating compatibility among *Bactrocera cucurbitae* (Diptera: Tephritidae) populations from three different origins. *Journal of Applied Entomology* 137(Issue s1): 69–74. <https://doi.org/10.1111/j.1439-0418.2010.01576.x>
- Stathopoulou, P. M., E. Asimakis, and G. Tsiamis. 2021. *Enterobacter*: one bacterium multiple functions, pp. 917–945. In J. Hendrichs, R. Pereira and M. J. B. Vreysen (eds.), *Area-wide integrated pest management. Development and field application*. CRC Press, Boca Raton, FL, USA.
- Steiner, E., W. Woods, D. O. McInnis, J. Lindsey, R. Fogliani, and R. Soopaya. 2013. Ginger root oil increases mating competitiveness of sterile Mediterranean fruit fly (Diptera: Tephritidae) in Western Australia. *Journal of Applied Entomology* 137(Issue s1): 103–112. <https://doi.org/10.1111/j.1439-0418.2011.01630.x>
- Tan, K. H., and R. Nishida. 1996. Sex pheromone and mating competition after methyl eugenol consumption in the *Bactrocera dorsalis* complex, pp. 147–153. In B. A. McPherson and G. J. Steck (eds.), *Fruit fly pests: a world assessment of their biology and management*. St. Lucie Press, Delray Beach, FL, USA.

- Tan, K. H., and R. Nishida. 2012.** Methyl eugenol: its occurrence, distribution, and role in nature, especially in relation to insect behavior and pollination. *Journal of Insect Science* 12: 56. <https://doi.org/10.1673/031.012.5601>
- Tan, L. T., and K. H. Tan. 2013.** Automated tephritid fruit fly semiochemical mass feeding structure: design, construction and testing. *Journal of Applied Entomology* 137(Issue s1): 217–229. <https://doi.org/10.1111/j.1439-0418.2011.01680.x>
- Taylor, P. W., and B. Yuval. 1999.** Postcopulatory sexual selection in Mediterranean fruit flies: advantages for large and protein-fed males. *Animal Behaviour* 58: 247–254. <https://doi.org/10.1006/anbe.1999.1137>
- Taylor, P. W., M. Khan, S. R. Collins, and O. L. Reynolds. 2013a.** Yeast hydrolysate supplement increases starvation vulnerability of Queensland fruit fly. *Physiological Entomology* 38: 337–343. <https://doi.org/10.1111/phen.12039>
- Taylor, P. W., D. Pérez-Staples, C. W. Weldon, S. R. Collins, B. G. Fanson, S. Yap, and C. Smallridge. 2013b.** Post-teneral nutrition as an influence on reproductive development, sexual performance and longevity of Queensland fruit flies. *Journal of Applied Entomology* 137(Issue s1): 113–125. <https://doi.org/10.1111/j.1439-0418.2011.01644.x>
- Teal, P. E. A., and Y. Gómez-Simuta. 2002.** Endogenous coordination on reproductive maturity and sexual signaling in tephritid fruit flies, pp. 469–474. *In* R. Keller, H. Dircksen, D. Sedlmeier and H. Vuadry (eds.), *Proceedings of the 21st Conference of European Comparative Endocrinologists*. Monduzzi Editore, Bonn, Germany.
- Teal, P. E. A., Y. Gómez-Simuta, and A. T. Proveaux. 2000.** Mating experience and juvenile hormone enhance sexual signaling and mating in male Caribbean fruit flies. *Proceedings of the National Academy of Sciences of the USA* 97(7): 3708–3712. <https://doi.org/10.1073/pnas.97.7.3708>
- Teal, P. E. A., Y. Gómez-Simuta, B. D. Deuben, T. C. Holler, and S. Olson. 2007.** Improving the efficacy of the sterile insect technique for fruit flies by incorporation of hormone and dietary supplements into adult holding protocols, pp. 163–173. *In* M. J. B. Vreysen, A. S. Robinson and J. Hendrichs (eds.), *Area wide control of insect pests. From research to field implementation*. Springer, Dordrecht, The Netherlands. <https://www.iaea.org/sites/default/files/area-wide-control-insect-pests-book.pdf>
- Teal, P. E. A., R. Pereira, D. F. Segura, I. Haq, Y. Gómez-Simuta, A. S. Robinson, and J. Hendrichs. 2013.** Methoprene and protein supplements accelerate reproductive development and improve mating success of male tephritid flies. *Journal of Applied Entomology* 137(Issue s1): 91–98. <https://doi.org/10.1111/j.1439-0418.2010.01606.x>
- Tween, G., and P. Rendón. 2007.** Current advances in the use of cryogenics and aerial navigation technologies for sterile insect delivery systems, pp. 229–238. *In* M. J. B. Vreysen, A. S. Robinson and J. Hendrichs (eds.), *Area-wide control of insect pests. From research to field implementation*. Springer, Dordrecht, The Netherlands. <https://www.iaea.org/sites/default/files/area-wide-control-insect-pests-book.pdf>
- (USDA/APHIS) United States Department of Agriculture/Animal and Plant Health Inspection Service. 2009.** Final Report: United States, Mexico, and Guatemala Fruit Fly Emergence and Release Facilities Review. Conducted July 2008. 81 pp. PDF document can be downloaded. Riverdale, MD, USA. https://www.researchgate.net/publication/240609561_United_States_Mexico_and_Guatemala_Fruit_Fly_Emergence_and_Release_Facilities_Review
- Utges, M. E., J. C. Vilardi, A. Oropeza, J. Toledo, and P. Liedo. 2013.** Pre-release diet effect on field survival and dispersal of *Anastrepha ludens* and *Anastrepha obliqua* (Diptera: Tephritidae). *Journal of Applied Entomology* 137(Issue s1): 163–177. <https://doi.org/10.1111/j.1439-0418.2011.01663.x>
- Vargas, R. I., and R. Prokopy. 2006.** Attraction and feeding responses of melon flies and oriental fruit flies (Diptera: Tephritidae) to various protein baits with and without toxicants. *Proceedings of the Hawaiian Entomological Society* 38: 49–60. https://scholarspace.manoa.hawaii.edu/bitstream/10125/178/5_vargas.pdf
- Vargas, R. I., N. W. Miller, and R. J. Prokopy. 2002.** Attraction and feeding responses of Mediterranean fruit fly and a natural enemy to protein baits laced with two novel toxins, phloxine B and spinosad. *Entomologia Experimentalis et Applicata* 102(3): 273–282. <https://doi.org/10.1046/j.1570-7458.2002.00948.x>
- Vargas, R. I., L. Leblanc, J. C. Piñero, and K. M. Hoffman. 2014.** Male annihilation, past, present, and future, pp. 493–511. *In* T. Shelly, N. Epsky, E. B. Jang, J. Reyes-Flores and R. I. Vargas (eds.),

- Trapping and the detection, control, and regulation of tephritid fruit flies: lures, area-wide programs, trade implications. Springer International Publishing, Dordrecht, The Netherlands.
- Vera, M. T., M. J. Ruiz, A. Oviedo, S. Abraham, M. Mendoza, D. F. Segura, N. A. Kouloussis, and E. Willink. 2013.** Fruit compounds affect male sexual success in the South American fruit fly, *Anastrepha fraterculus* (Diptera: Tephritidae). *Journal of Applied Entomology* 137(Issue s1): 2–10. <https://doi.org/10.1111/j.1439-0418.2010.01516.x>
- Wee, S. L., and K. H. Tan. 2001.** Allomonal and hepatotoxic effects following methyl eugenol consumption in *Bactrocera papayae* male against *Gekko monarchus*. *Journal of Chemical Ecology* 27: 953–964. <https://doi.org/10.1023/A:1010387020135>
- Wee, S. L., K. H. Tan, and R. Nishida. 2007.** Pharmacophagy of methyl eugenol by males enhances sexual selection of *Bactrocera carambolae*. *Journal of Chemical Ecology* 33: 1272–1282. <https://doi.org/10.1007/s10886-007-9295-0>
- Wee, S. L., M. Z. Abdul Munir, and A. K. W. Hee. 2018.** Attraction and consumption of methyl eugenol by male *Bactrocera umbrosa* Fabricius (Diptera: Tephritidae) promotes conspecific sexual communication and mating performance. *Bulletin of Entomological Research* 108: 116–124. <https://doi.org/10.1017/S0007485317000554>
- Weldon, C. W., and P. W. Taylor. 2011.** Sexual development of wild and mass-reared male Queensland fruit flies in response to natural food sources. *Entomologia Experimentalis et Applicata* 139: 17–24. <https://doi.org/10.1111/j.1570-7458.2011.01104.x>
- Weldon, C. W., D. Perez-Staples, and P. W. Taylor. 2008.** Feeding on yeast hydrolysate enhances attraction to cue-lure in Queensland fruit flies, *Bactrocera tryoni*. *Entomologia Experimentalis et Applicata* 129: 200–209. <https://doi.org/10.1111/j.1570-7458.2008.00768.x>
- Woruba, D. N., J. L. Morrow, O. L. Reynolds, T. A. Chapman, D. P. Collins, and M. Riegler. 2019.** Diet and irradiation effects on the bacterial community composition and structure in the gut of domesticated teneral and mature Queensland fruit fly, *Bactrocera tryoni* (Diptera: Tephritidae). *BMC Microbiology* 19: 281. <https://doi.org/10.1186/s12866-019-1649-6>
- Yao, M., H. Zhang, P. Cai, X. Gu, D. Wang, and Q. Ji. 2017.** Enhanced fitness of a *Bactrocera cucurbitae* genetic sexing strain based on the addition of gut-isolated probiotics (*Enterobacter* spec.) to the larval diet. *Entomologia Experimentalis et Applicata* 162: 197–203. <https://doi.org/10.1111/eea.12529>
- Yuval, B., R. Kaspi, S. A. Field, S. Blay, and P. Taylor. 2002.** Effects of post-teneral nutrition on reproductive success of male Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* 85: 165–170. [https://doi.org/10.1653/0015-4040\(2002\)085\[0165:EOPTNO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2002)085[0165:EOPTNO]2.0.CO;2)
- Yuval, B., M. Maor, K. Levy, R. Kaspi, P. Taylor, and T. Shelly. 2007.** Breakfast of champions or kiss of death? Survival and sexual performance of protein-fed, sterile Mediterranean fruit flies (Diptera: Tephritidae). *Florida Entomologist* 90: 115–122. [https://doi.org/10.1653/0015-4040\(2007\)90\[115:BOCOKO\]2.0.CO;2](https://doi.org/10.1653/0015-4040(2007)90[115:BOCOKO]2.0.CO;2)
- Yuval, B., E. Ben-Ami, A. Behar, M. Ben-Yosef, and E. Jurkevitch. 2013.** The Mediterranean fruit fly and its bacteria – potential for improving sterile insect technique operations. *Journal of Applied Entomology* 137(Issue s1): 39–42. <https://doi.org/10.1111/j.1439-0418.2010.01555.x>