80 Viral hepatitis in pregnancy

Dennis Ng

This chapter reviews two of the most common hepatitis affecting pregnancy, hepatitis B and C. It is catered toward a mixed audience of practicing obstetricians, obstetrics and gynecology residents, maternal–fetal medicine fellows, and health-care providers for pregnant patients. There has been an explosion of knowledge in disease pathophysiology and treatment options for hepatitis B and C in the last 10 years. Guidelines for treatment and management for hepatitis B and C are changing every 2 to 3 years. To the uninitiated, this can be a daunting subject.

My intention is to provide you with an easy-to-understand review of the subject, emphasizing on the main concepts and principles in the management of hepatitis B and C, in the context of pregnancy and delivery. As such, no attempt has been made nor is it possible to be all-inclusive in the minute details of the subject. Instead, I attempt to provide you with an insight into the pertinent questions at hand related to pregnancy and delivery that may arise in your consultation. Particular emphasis will be placed on counseling and public health issues, a subject often forgotten by practitioners, but of utmost importance.

Interested readers may refer to the several excellent textbooks on hepatology, review articles, and guidelines published by American Association for the Study of Liver Diseases (AASLD). This chapter will focus only on hepatitis B and C. Brief mention of hepatitis E will be made. Other viral hepatitis like herpes simplex virus hepatitis will not be covered.

PRESENTATION OF HEPATITIS B AND C AND GENERAL APPROACH

The pregnant female may acquire hepatitis A, B, and C acutely. If symptomatic, they usually complain of flu-like symptoms, general aches and pains, fatigue, nausea, and vomiting. Physical exam may reveal icterus, jaundice, right upper quadrant tenderness, and change in the color of urine. Labs may reveal high levels of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) in the thousand range. alkaline phosphatase is usually only slightly elevated. Bilirubin may be elevated. These pregnant patients are best admitted to the hospital, particularly if symptomatic with nausea, vomiting, dehydration, where an expedited workup can be performed to rule out other causes of liver diseases unique to pregnancy like acute fatty liver of pregnancy and hemolysis, elevated liver enzymes, and low platelet count (HELLP) syndrome.

More commonly, hepatitis B comes to the attention of the clinician from the adoption of the universal hepatitis B screening during prenatal visit in the United States. Hepatitis C should be tested in high-risk patients. If found to be positive, they should be referred to the hepatologist/gastroenterologist to determine the status of hepatitis B or C, severity of liver

disease, and to help with management of liver disease postdelivery. It is important to screen for these diseases because many times, for a young healthy female patient, the obstetrician may be the only doctor they will see. Once diagnosed, it allows the clinician the golden opportunity to talk about public health issues, mode of transmission of hepatitis, ways to prevent progression of liver disease (e.g., quit drugs and alcohol), referral to a hepatologist for possible treatment options, and cancer screening for the liver (if relevant).

HEPATITIS B IN PREGNANCY

Epidemiology

Worldwide more than 350 million individuals are chronic hepatitis B carriers [1]. The prevalence of hepatitis B varies from region to region. Hepatitis B prevalence can be divided into low-prevalence (e.g., North America, Western Europe), intermediate-prevalence, and high-prevalence areas (e.g., sub-Saharan Africa, Asia excluding Japan, the Pacific basin, parts of the Middle East, and the Amazon basin).

It is important to note that there is a recent increase in immigrants from highly prevalent countries over the last few decades into the United States. Between 1994 and 2003, the United States admitted about 40,000 individuals with chronic hepatitis B infection annually [2]. Hence, areas of high concentration of these immigrants will have a higher prevalence rate. For example, a recent study in 2009 examined the seroprevalence of HepBsAg among the Asian Chinese in an urban Chinatown internal medicine practice in Chicago [3]. The number was 11.1%, suggesting that we should be even more vigilant about screening high-risk patients, because treatment where indicated has been shown to improve prognosis and decrease morbidity and mortality from hepatitis B in the long run.

Acute hepatitis B occurs in 1 to 2 out of 1000 pregnancies in the United States. Chronic carrier state occurs in 6 to 10 of 1000 pregnancies.

Importance of Hepatitis B on a Global Scheme

It is generally accepted that at least 50% of the hepatitis B carriers acquire infection either perinatally or during early childhood. Studies suggest that most of the infections occur around the time of delivery. This gives us the unique opportunity of intervening by providing active and passive hepatitis B vaccination. The goal is to prevent infection in the newborn, hence decreasing the burden of hepatitis B infection worldwide.

When you acquire hepatitis B as an adult, you have a 90% to 95% chance of clearing this virus (resolved hepatitis B infection). But when you acquire this as an infant, you have only a 5% to 10% chance of clearing this virus, that is, most of

the infants will be chronic carriers of hepatitis B, with its disease-related morbidity and mortality. Therefore, it makes sense for us to intervene to decrease the rate of perinatal and childhood infection. Taiwan has implemented a mandatory hepatitis B vaccination program 20 years ago, with a significant impact on perinatal, childhood, and adolescent infection. Chronic liver disease and hepatocellular carcinoma attributed to hepatitis B have decreased since implementation [4].

In the United States, we have a mandatory screening for all pregnant females, and if positive for hepatitis B surface antigen, the infants are to get active and passive immunization.

Even in the best of hands, when vaccinations have been carried out appropriately, there is still about 10% of infants who will get hepatitis B infection, suggesting that perhaps they acquired this virus in utero [5,6].

Effect of Acute Hepatitis B Infection During Pregnancy

Acute hepatitis B can occur during pregnancy; they present the same way as in the general population. Acute hepatitis B must be differentiated from hepatitis A and C, and other causes of hepatitis unique to pregnancy (HELLP syndrome, acute fatty liver of pregnancy, etc.). It appears that acute hepatitis B does not increase mortality and has no teratogenic side effects. However, a higher incidence of low birth weight and prematurity has been reported [7].

Effect of Pregnancy on Chronic Hepatitis B

In general, patients do well during pregnancy. Patients have been known to have a flare of hepatitis B during or after delivery (one theory is that the high levels of adrenocorticosteroids in the body have now decreased significantly postpartum, just like abruptly removing steroids from the patient). Therefore, it is important to monitor patients for symptoms of hepatitis and also to test for liver function tests, during and also after delivery.

Diagnosing Hepatitis B in the Mother

Over the years, we have developed better tests for hepatitis B, including the ability to measure HBV DNA PCR quantitatively down to the most minute level, for example, 5 IU/mL.

The most common mistake that I see clinicians make is to order an acute hepatitis panel. Usually this consists of HepBsAg, HepBcIgM, HepCAb, and HepAIgM. These are acute hepatitis markers, meaning good for diagnosing acute hepatitis infections, but tell you nothing about the chronic state of hepatitis B, which is what we are interested in, to determine treatment and management for the pregnant patient.

Below Are Some Typical Hepatitis B Serology You Might Encounter in Clinical Practice

Case 1

A 20-year-old Vietnamese pregnant college student, sent to you because tested positive for hepatitis B. Further workup reveals liver function tests (LFTs) normal, HepBsAg positive, HepBsAb negative, HepBcTotal positive, HepBcIgM negative, HepBeAg positive, HepBeAb negative, and HBV DNA >2,000,000,000. The ultrasound (US) of liver unremarkable.

Impression: She is in the immunotolerant phase of hepatitis B.

Case 2

A 30-year-old Malaysian female, pregnant, family history positive for hepatitis B in mother and father. Further workup reveals LFTs elevated, AST 60, ALT 70, HepBsAg positive, HepBsAb negative, HepBcTotal positive, HepBcIgM negative, HepBeAg positive, HepBeAb negative, HBV DNA 200,000. The US of liver with echogenic liver.

Impression: She is in immunoclearance phase of hepatitis B.

Case 3

A 38-year-old Chinese female, pregnant, family history positive of liver cancer in her dad. Further workup reveals LFTs normal, HepBsAg negative, HepBsAb positive, HepBcTotal positive, HepBcIgM negative, HepBeAg negative, HepBeAb positive, and HBV DNA undetectable. The US of liver unremarkable.

Impression: She has resolved hepatitis B infection.

Case 4

A 22-year-old white American female with the following labs: LFTs normal, HepBsAb positive, and every other marker for hepatitis B negative.

Impression: She has been vaccinated in the past and has shown immunity as evidenced by antibody formation.

Vaccinations for Hepatitis B for the Mother and Baby

The Centers for Disease Control and Prevention (CDC) recommends that all mothers be tested for hepatitis B during the initial prenatal visit [8]. This applies even if the patient has been vaccinated in the past or tested in the previous pregnancy.

If negative, the pregnant mother who is at risk for hepatitis B infection (unsafe sex, multiple sexual partners, health-care workers, active intravenous drug user, etc.) should be vaccinated. The limited data we have indicate that there is no apparent risk for adverse events to developing fetuses. Current vaccines contain noninfectious HBsAg and should cause no risk to the fetus.

If HepBsAg is found to be negative, and the high-risk patient refuses vaccination, or is unable or unwilling to get vaccinated, then HepBsAg should be rechecked closer to the time of delivery because this will change management if found to be positive.

If HepBsAg positive, the patient is a carrier for hepatitis B. She should be referred to a gastroenterologist/hepatologist for further characterization of her liver disease and the need for treatment. Counseling should be provided regarding the risk of transmission to the newborn and sexual partners, and prevention of HBV transmission to newborns and sexual partners. They should be evaluated for other concomitant liver diseases including hepatitis A and hepatitis C. All sexual contacts and family members should be tested for hepatitis B. If the patient is not immune to hepatitis A and is at high risk of acquiring it, then vaccines for hepatitis A should be given, because of the risk of severe hepatitis with hepatitis A on top of hepatitis B infection [9].

A patient who is not a hepatitis B treatment candidate during pregnancy may be a candidate for treatment at a later point in time. Therefore, it should be emphasized to the patient that she should continue to follow up with the

gastroenterologist/hepatologist after delivery. Hepatitis B is a chronic disease that requires regular monitoring and liver cancer screening in at risk patients. The clinicians caring for these pregnant patients have the golden opportunity of providing education regarding hepatitis B and also to ensure that patients continue to follow up with specialists.

The pregnant patient with hepatitis B should be enrolled into a special program that ensures that active and passive vaccinations not only happen, but also happen in a timely manner. This includes making sure that there is ready access to the vaccines when needed, including at 3:00 in the morning or the weekend. We have heard of horror stories in smaller hospitals where the vaccines is available but locked up in the central pharmacy in the weekend. This program should also ensure that the hepatitis B vaccines are completed in a timely fashion postdelivery.

Treatments Available for Hepatitis B in 2010

The last 10 years have seen tremendous progress in the treatment of hepatitis B. We now have five FDA-approved nucleoside and nucleotide analogs and two interferon-based regimen:

Interferonalpha-2b recombinant (1992), lamivudine (1998), adefovir dipivoxil (2002), entecavir (2005), peginterferon alpha-2a (2005), telbivudine (2006), and tenofovir (2008).

It is beyond the scope of this chapter to discuss selection of hepatitis B drugs and the rationale behind this. For the interested readers, please refer to AASLD 2009 updated guidelines.

Tenofovir and telbivudine are pregnancy category class B, while the rests are class C.

Prevention of Hepatitis B Transmission to Newborns

Active and Passive Vaccination

CDC recommendation for hepatitis B in infants is stated below [8]:

At Birth

- Administer monovalent hep B to all newborns before hospital discharge.
- If mother is HBsAg positive, administer HepB and 0.5mL of hepatitis B immunoglobulin (HBIG) within 12 hours of birth.
- If mother is HBsAg unknown, administer HepB within 12 hours of birth. Determine mother's HBsAg status as soon as possible and, if HBsAg positive, administer HBIG no later than age 1 week.

After the Birth Dose

- The HepB series should be completed with either monovalent HepB or a combination vaccine containing HepB. The second dose should be administered at age 1 or 2 months. Monovalent HepB vaccine should be used for doses administered before age 6 weeks. The final dose should be administered no earlier than age 24 weeks.
- Infants born to HBsAg-positive mothers should be tested for HBsAg and antibody to HBsAg 1 to

2 months after completion of at least three doses of the HepB series, at age 9 through 18 months.

- Administration of four doses of HepB to infants is permissible when a combination vaccine containing HepB is administered after the birth dose. The fourth dose should be administered no earlier than age 24 weeks.

Breast-Feeding in Hepatitis B Patients

Even though HepBsAg can be detected in breast milk in a large proportion of infected women, several studies fail to demonstrate increased risk of infection due to breast-feeding as opposed to infants who are formula-fed in hepatitis B-infected mother.

Breast-feeding is not considered a contraindication to breast-feeding by the CDC or American Academy of Pediatrics in infants who receive HBIG and active vaccination as indicated.

The nipples should be cared for properly, and mothers should not breast-feed if there are signs of cracked or bleeding nipple.

Mode of Delivery

Despite the attractive idea of decreasing the risk of perinatal transmission, studies have so far shown that the mode of delivery does not appear to have a significant effect on decreasing the risk of hepatitis B transmission [10]. The CDC or American College of Obstetricians and Gynecologists does not currently recommend caesarian section as the preferred mode of delivery purely for the purpose of reducing perinatal transmission.

Role of Antiviral

There is emerging evidence that patients who are put on antiviral therapy during the last trimester, to reduce hepatitis B viral load, could potentially reduce the risk of perinatal transmission. There are many factors to be taken into consideration, including compliance with therapy, severity of liver disease, history of prior treatment, resistance issues, the risk of flare of hepatitis B should the antiviral be stopped postpartum, mothers' preference, and so on. The major concern is that three of the medications we have for hepatitis B are in pregnancy category C and two in category B. There is no category A medication for hepatitis at this point in time. These decisions are best made by the hepatologist and patient together.

Summary for Hepatitis B

Hepatitis B when diagnosed in mother prior to delivery affords us the opportunity to make a difference in reducing the risk of mother-to-infant transmission. Besides vaccination of the newborn, other important aspects of care of the young pregnant patients include counseling regarding their liver disease and mode of transmission and ways to reduce the risk of progression of liver disease. Patients who are not candidates for treatment now may become candidates in the future. Hepatitis B is not a disease to be diagnosed at a single point in time and be forgotten about. It requires lifelong follow-up with a clinician who is familiar with this disease.

HEPATITIS C

Background

Hepatitis C is a major public health problem in the United States. It is estimated that 170 million people are infected worldwide [11]. This is the number one cause for liver transplantation in the United States today. The top three diagnoses in my liver clinic are hepatitis C, nonalcoholic steatohepatitis, and alcoholic liver disease. The sad part is that the majority of the patients who are diagnosed with hepatitis C are unaware of their liver disease, because hepatitis C mostly do not present with any specific symptoms. In fact, most patients will tell you that they do not feel any different from their normal healthy self. Therefore, it is even more important to screen high-risk patients for active disease.

Why Do We Worry About Testing for Hepatitis C in High-Risk Patients?

We worry because about 80% to 90% of patients who acquire acute hepatitis C infection fail to clear the virus. After 6 months from the time of exposure, if they still have virus in their system, they are considered to have chronic hepatitis C with active disease. Over time, this may lead to increase in fibrosis of the liver, progression of liver disease to cirrhosis, and finally, decompensated cirrhosis (ascites, edema, hepatic encephalopathy, etc.), and death. This is in contradistinction to hepatitis B, where if they get infected as an adult, about 90% to 95% of patients successfully clear the virus (resolved hepatitis B infection).

Practicing obstetricians and health-care providers during and after pregnancy have a unique opportunity to screen high-risk patients for hepatitis C. Many times, this may be the only time when young ladies touch base with the health-care system. They may not have a primary care provider. This is even more true in patients who are poor, have no insurance, and where drug and alcohol usage is a daily occurrence.

Who Are High-Risk Patients?

The AASLD 2009 guidelines suggest that the following people are high risk for having hepatitis C infection and should be screened for hepatitis C [12]:

1. persons who have injected illicit drugs in the recent and remote past, including those who injected only once and do not consider themselves to be drug users,
2. persons with conditions associated with a high prevalence of HCV infection including
 - persons with HIV infection, persons with hemophilia who received clotting factor concentrates prior to 1987, persons who have ever been on hemodialysis, and persons with unexplained abnormal LFTs,
3. prior recipients of transfusions or organ transplants prior to July 1992 including
 - persons who were notified that they had received blood from a donor who later tested positive for HCV infection, persons who received a transfusion of blood or blood products, and persons who received an organ transplant,
4. children born to HCV-infected mothers,

5. health-care, emergency medical, and public safety workers after a needlestick injury or mucosal exposure to HCV-positive blood, and
6. current sexual partners of HCV-infected persons

How Do We Make the Diagnosis?

Over the last 10 years, we have made a remarkable improvement in the diagnostic sensitivity of assays to detect hepatitis C RNA virus. We can now detect virus level as low as 5 IU/mL.

For high-risk patients, screening for chronic hepatitis C should begin with anti-HCV ELISA test (using the latest generation). A positive result suggests exposure to the virus in the past. The next test that should be done is HCV RNA PCR and a genotype. If PCR is positive, this implies that the patient has active chronic hepatitis C and should be referred to the appropriate specialist for consideration of treatment and education.

What Treatments Are Available Now?

Hepatitis C treatment has come a long way since it was discovered. The perception in the past was that you cannot cure hepatitis C. Now we have effective treatments, but unfortunately, cure rate is not 100%. On average, the quoted sustained virologic response (SVR; we think of this as cure) rate is about 50%, but this is an oversimplistic view. Many factors affect the SVR, including gender, race, obesity, concurrent liver disease, severity of liver disease, coinfection with HIV, on-treatment viral kinetics, genotype, compliance with therapy, and so on. The latest STAT-C drugs that are projected to come out in 2011 look promising in increasing the SVR.

What Does the Treatment Entail?

It involves a subcutaneous shot (peginterferon) every week and ribavirin pills twice a day. Treatment duration depends on prior treatment response, genotype of virus, viral kinetics during response, and so on and can range anywhere from 6 months to 12 months to 18 months. This will be determined by the specialist based on prior treatment response, on-treatment response, and other factors. Treatment may also be terminated early because of development of intolerable side effects, or development of life-threatening side effects, or if the patient is deemed a nonresponder to interferon therapy.

Who Are Good Candidates for Treatment?

The AASLD guidelines suggest that every patient who has chronic hepatitis C should be considered treatment candidates, unless there is contraindication. This includes patients in prisons and those who are on methadone therapy. We know that treatment can lead to SVR (i.e., the HCV RNA remains undetectable at 6 months after stopping therapy, i.e., cured). This will prevent the progression of liver disease to cirrhosis, translating into a decrease in morbidity and mortality.

Unfortunately, this therapy is neither suitable nor possible for everyone because of the severe and sometimes life-threatening side effects of interferon and ribavirin therapy. In addition, patients with certain preexisting medical conditions are not suitable candidates for treatment. Cost can be an issue. Also treatment requires the ability and willingness to take the medications faithfully, because SVR will be affected by

lack of compliance. They need to be able to show up regularly for appointments for monitoring.

Treatment with Regard to Pregnant Patients?

Hepatitis C treatment is contraindicated during pregnancy. Pegylated interferon is a pregnancy category C drug, and ribavirin is a pregnancy category X drug. It is important that the patient be referred to see a specialist so that they can be followed up postpartum for possible treatment in the future.

In addition, she should be educated on the mode of transmission for hepatitis C and abstinence from alcohol, tested for other sexually transmitted diseases, and advised to get her partners tested [12].

During hepatitis C treatment, patients are not allowed to get pregnant. It is recommended that they use two forms of contraception. I do not treat any patients who are unable or unwilling to comply with this. The package insert for pegylated interferon recommends that patients do not get pregnant until 6 months post therapy. Practically speaking, if patients are very worried about their hepatitis C, and would like to get treated first before getting pregnant, this will usually mean a delay of 1 to 2 years depending on how they respond to therapy.

Another frequently asked question is whether they should get treated first before pregnancy? This will depend on the age of the patient, the severity of liver disease, patient's preference, and urgency of getting pregnant and is best discussed with the specialist.

It is important to note that treatment with interferon can cause or exacerbate preexisting depression. In addition, treatment is contraindicated when she develops postpartum depression.

Please bear in mind that patients who are not treatment candidates now may become suitable candidates later in time and, therefore, the importance of referring to specialists for follow-up.

Transmission of Hepatitis C to Infants

Hepatitis C is primarily transmitted by parenteral exposure to blood and blood products from HCV-infected persons. This used to be the most common cause of hepatitis C infection in infants and children, until we developed ways to screen for hepatitis C in blood. Now, vertical transmission is mainly responsible for infection in newborn infants. It is important to identify these infants because the majority of these patients will remain chronically infected with hepatitis C, hence the potential to develop chronic liver disease.

The rate of vertical transmission of hepatitis C is estimated to be 5%. This number is reported to be higher in the 10% to 15 % range in HIV–hepatitis C coinfecting patients [13].

Numerous studies have suggested that the vertical risk of transmission for hepatitis C is increased in the following setting:

1. HIV–hepatitis C coinfection.
2. HCV viremia—mothers with positive virus load have increased risk of transmission, compared with mothers with undetectable viral load (suggesting resolved infection). The overwhelming majority of infants with HCV infection have mothers with positive hepatitis C virus in the blood.

3. Prolonged rupture of membrane for more than 6 hours has been associated with increased risk of perinatal transmission.
4. Obstetric procedures and intrapartum exposure to maternal blood infected by HCV—Steininger et al. have shown that invasive monitoring of fetus with scalp electrode increases the risk of transmission [14].

What Can We Do to Decrease the Risk of Transmission to the Infant?

Unlike hepatitis B, no effective intervention has been identified so far to decrease the risk of transmission of HCV to infants. There have been many studies looking to see if C-section will decrease the risk compared with vaginal delivery. Most of the studies suggest that the risk of perinatal transmission is not altered. The Cochrane Collaboration (Issue 2 2009) has published a review on this topic [15]. The authors suggest that there is no good evidence to support using C-section for reducing mother-to-baby transmission of hepatitis C during labor and birth.

To the mother-to-be who is absolutely horrified by the idea of hepatitis C transmission to their newborn, a case could be made to treat their hepatitis C first prior to getting pregnant, because we know that the risk of hepatitis C transmission is much less in a patient who has no detectable virus in the blood versus active hepatitis C infection with high viral load. They must be made to understand that this will delay pregnancy anywhere from 1 to 2 years (see above discussion).

Testing for Hepatitis C in Infants

The antibodies from the mother have been found to be in the infants even though the infants are not infected; therefore, finding positive HCV antibodies in the newly born infant means very little. Antibodies have been tested positive in infants as late as 15 months old [16].

The NIH consensus statement in 2002 suggested the following for diagnosis of perinatal infections [17]:

1. infants born to HCV-positive mothers should be tested for HCV infection by HCV RNA tests on two occasions between the ages of 2 and 6 months and/or
2. have tests for anti-HCV after 15 months of age.

Breast-Feeding in Hepatitis C Patients

Hepatitis C has been found in low levels in breast milk of HCV RNA–positive mothers. Multiple studies have not shown an increased risk of vertical transmission via breast-feeding.

Large cohorts of mothers infected with HCV suggest that in the absence of damaged, cracked, or bleeding nipples, breast-feeding can be performed safely without increasing the risk of transmission to the infant.

CDC currently (October 2009) suggests that it is safe for mother infected with HCV to breast-feed unless she has a cracked or bleeding nipple.

Summary

Hepatitis C is a common disease in high-risk patients. They should be screened for hepatitis C, even though there is no

known intervention to decrease the risk of hepatitis C from mother to infant. Pregnant patients are not candidates for treatment with interferon and ribavirin. They should be referred to specialist to determine whether they may be candidates for hepatitis C treatment in the future. Education should be provided regarding public health–related issues, including mode of transmission of hepatitis C, avoidance of alcohol, vaccinations for hepatitis A and B if appropriate, and testing of sexual partners. It is fine to breast-feed the infant unless the nipples are cracked or bleeding.

HEPATITIS E

Clinically apparent acute hepatitis E is uncommon in the United States. A brief mention will be made here, because of its severity when acquired during pregnancy.

Hepatitis E causes acute hepatitis, just like hepatitis A. It is predominantly transmitted via the fecal–oral route, but less frequent mode of transmission has also been reported (ingestion of raw or undercooked shellfish, infrequently by blood products or blood transfusion).

Hepatitis E usually causes a self-limited acute hepatitis in men and nonpregnant females. However, HEV in pregnant females can result in fulminant hepatic failure, especially in the last trimester, resulting in death in as high as 25% of patients [18]. This is especially prominent in particular geographic areas in India.

This infection should be in the differential diagnosis of any female patient with a travel history to endemic areas, who presents with acute hepatitis. Vaccines for hepatitis E are not yet commercially available. Treatment remains supportive. It is best to educate pregnant patients who will travel to endemic areas (i.e., Asia, Africa, Middle East, and Central America) to avoid uncooked shellfish, avoid water that is not boiled, and avoid uncooked fruits and vegetables.

ADDENDUM

Since the writing of this chapter, 2 new drugs (telaprevir and boceprevir) have been approved by the FDA for the treatment of hepatitis C genotype 1. They significantly improve the cure rate for hepatitis C treatment. However, these protease inhibitors must be used together with peg-interferon and ribavirin, which means that they cannot be used for pregnant patients. All the discussion for the treatment of hepatitis C in pregnant patients still applies. Those interested to read more may refer to the AASLD 2011 Hepatitis C Treatment Guidelines.

ACKNOWLEDGMENTS

I would like to acknowledge the help of Jessica Wynn, MD, Internal Medicine Resident PGY2, University of Missouri, Columbia, Missouri, in preparing the section on *hepatitis E* in this chapter.

REFERENCES

1. Maynard JE. Hepatitis B: global importance and need for control. *Vaccine* 1990; 8(Suppl): S18–20.
2. Wasley A, Grytdal S, Gallagher K. Surveillance for acute viral hepatitis—United States, 2006. *MMWR Surveill Summ* 2008; 57: 1–24.
3. Cotler SJ, Dhamija MK, Luc BJ, et al. The prevalence and clinical correlates of elevated ALT levels in an urban Chinatown community. *J Viral Hepat* 2010; 17: 148–52.
4. Chan CY, Lee SD, Lo KJ. Legend of hepatitis B vaccination: the Taiwan experience. *J Gastroenterol Hepatol* 2004; 19: 121–6.
5. Zhu Q, Lu Q, Gu X, Xu H, Duan S. A preliminary study on interruption of HBV transmission in uterus. *Chin Med J (Engl)* 1997; 110: 145–7.
6. Stevens CE, Toy PT, Tong MJ, et al. Perinatal hepatitis B virus transmission in the United States. Prevention by passive-active immunization. *JAMA* 1985; 253: 1740–5.
7. Jonas MM. Hepatitis B and pregnancy: an underestimated issue. *Liver Int* 2009; 29(Suppl 1): 133–9.
8. Mast EE, Margolis HS, Fiore AE, et al. A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP) part 1: immunization of infants, children, and adolescents. *MMWR Recomm Rep* 2005; 54: 1–31. Erratum in: *MMWR Morb Mortal Wkly Rep*. 2006 Feb 17; 55(6): 158–9. *MMWR Morb Mortal Wkly Rep*. 2007 Dec 7; 56(48): 1267.
9. Lok AS, McMahon BJ. Chronic hepatitis B: update 2009. *Hepatology* 2009; 50: 661–2.
10. Yang J, Zeng XM, Men YL, Zhao LS. Elective caesarean section versus vaginal delivery for preventing mother to child transmission of hepatitis B virus—a systematic review. *Virol J* 2008; 5: 100.
11. World Health Organization, “Hepatitis C. Epidemic and Pandemic Alert and Response,” 2002. [Available from: <http://www.who.int/csr/disease/hepatitis/whocdscrlyo2003/en/index.html>.].
12. Ghany MG, Strader DB, Thomas DL, et al. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology* 2009; 49: 1335–74.
13. Su GL. itis C in pregnancy. *Curr Gastroenterol Rep* 2005; 7: 45–9.
14. Steininger C, Kundi M, Jatzko G, et al. Increased risk of mother-to-infant transmission of hepatitis C virus by intrapartum infantile exposure to maternal blood. *J Infect Dis* 2003; 187: 345–51.
15. McIntyre PG, Tosh K, McGuire W. Caesarean section versus vaginal delivery for preventing mother to infant hepatitis C virus transmission. *Cochrane Database Syst Rev* 2006; CD005546.
16. Indolfi G, Resti M. Fetal transmission of hepatitis C virus infection. *J Med Virol Fetal* 2009; 81: 836–43.
17. Boyer JL, et al. NIH Consensus Statement on Management of Hepatitis C: 2002. *NIH Consens State Sci Statements* 2002; 19: 1–46.
18. Khuroo MS, Teli MR, Skidmore S, et al. Incidence and severity of viral hepatitis in pregnancy. *Am J Med* 1981; 70: 252–5.