81 Toxoplasmosis

Edward R. Newton

INTRODUCTION

Although toxoplasma encephalitis (1) has gained worldwide recognition as a major cause of death and morbidity in immunodeficient individuals, especially those with acquired immunodeficiency syndrome (AIDS), the fetus and child remain the major victims of toxoplasmic infection. Approximately 4000 pregnant women per year in the United States are infected with *Toxoplasma gondii* and 10% to 60% of their fetuses will be infected. Although the majority (>90%) appear asymptomatic at birth, long-term studies suggest that a significant proportion (20–40%) will demonstrate defects consistent with chronic toxoplasmosis. These defects include chorioretinitis (most common), hydrocephalus, deafness, and mental retardation. Unfortunately, problems in primary prevention, diagnosis, treatment, and follow-up limit the effectiveness of intervention programs.

Biology

T. gondii is an ubiquitous coccidian whose effects are felt in many animals in addition to humans (Table 1) (2). Toxoplasmosis is a member of the phylum Apicomplexa, similar to malaria and cryptosporidia. Much of its life cycle is spent as an obligate intracellular parasite. There appear to be three distinct clonal genotypes with different human virulence; type II accounts for the majority of maternal to fetal infections (85–95%); 5% to 15% are caused by clonal type I.

The organism has three distinct forms: sporozoites within an oocyst, and tachyzoites (also called endozoites or trophozoites) and bradyzoites (or cystozoites) within a tissue cyst. The life cycle of *T. gondii* is outlined in Figure 1. Members of the cat family, Felidae, are the only known definitive hosts. The sexual reproductive cycle occurs in the villi of the cat ileum. The time from primary inoculation to oocyst appearance in the feces varies by the mode of primary infection: ingestion of tissue cysts (chronic infection in prey animals), 3 to 5 days; ingestion of fecal oocysts, 20 to 24 days; or ingestion of tachyzoites, 7 to 10 days. Oocysts are shed for 1 to 3 weeks with peak production between 5 and 8 days (4,5). Millions of oocysts are shed in a single day. Most cats are infected primarily between 6 and 12 weeks of age (6), corresponding to weaning and the development of hunting skills. Once a cat has been infected, its immune system severely curtails or eliminates future oocyst production. The previously infected cat may be a safer pet than curtailing the behavior of a seronegative cat.

After excretion with feces, the zygote within the oocysts ($13 \times 11 \mu\text{m}$) undergoes sporulation. The zygote divides into two sporoblasts that form two sporocysts within the larger oocyst. Each sporoblast undergoes two divisions to produce four sporozoites ($8 \times 3 \mu\text{m}$). Ultimately each oocyst contains eight infectious sporozoites. Sporulation takes 1 to 21 days

(average, 3–4 days), depending upon temperature, moisture, and oxygenation. The oocysts can remain infectious for months (4,5).

After elimination and within the 2 to 4 days prior to sporulation, the fecal oocysts are not infectious (4,5,7). As a result, human infection from direct contact with a shedding cat is unusual. The major role of cats in human toxoplasmosis is to pollute transiently the environment with massive numbers of oocysts that may remain infectious for months or years. A small coastal community in California contained 7284 owned cats and 2046 feral cats. Forty-four percent of the owned cats defecated outside >75% of the time. The annual fecal deposition of owned cats amounted to 76.4 tons (wet weight); the feral cats contributed about 29.5 tons of feces annually to the environment (8). In one estuarine in Northern California, degradation of the wetlands and runoff from nearby communities may have played a major role in the die-off of sea otters from a toxoplasmosis epidemic (9). Global warming and environmental degradation may increase the incidence of human infections with toxoplasmosis in northern Europe (10, 11). In northern Poland, about 18% of sand pits or sand heaps in three cities contained toxoplasma DNA, which was identified by PCR (12).

The potential public health implications are concerning. The likelihood of human toxoplasmosis is related to environmental conditions that enhance or reduce oocyst survival, the prevalence of feline toxoplasmosis within that region, and culinary habits of the at-risk population. There have been several studies looking at sources of toxoplasma infection in pregnant women (13–15). All three studies demonstrated the same risk factors for toxoplasmosis infection: consumption of under-cooked meat or game, contact with soil (poor hand hygiene), consumption of poorly cleansed raw vegetables (poor kitchen hygiene), poor cleansing of meat preparation utensils or surfaces (poor kitchen hygiene), and changing the litter box of the pet cat. The odds ratio exceeded 3.0 with all risk factors. Consumption of undercooked meats (50%), soil exposure (18%), and work with animals (5%) accounted for the major of maternal to fetal transmission of toxoplasma.

Higher rates of seropositivity to toxoplasmosis among pregnant women are associated with warm, moist climates: central Africa, 81%; Lisbon Portugal, 64%; Riyadh and Saudi Arabia, 30%; western Scotland, 13%; and Norway, 10% (16). In the United States, the age-adjusted seroprevalence rate was northeast, 29%; south, 23%; Midwest, 21%; and West, 18% (3). These observations are supported by experimental data concerning oocyst survival related to humidity and temperature. Oocysts remain infectious for at least 17 months in water, at least 32 days in 100% relative humidity (at 22–26°C), and 11 days at 37% relative humidity and lose their ability to infect at less than 19% relative humidity and when the moisture content of feces drops to 0% to 1% (17,18). Ambient

temperatures affect both sporulation (infectivity) and survival of the oocysts. Unless combined with low relative humidity (<20%), extremes in environmental temperature will not completely eliminate *T. gondii* from contaminated soil.

Survival and infectivity of sporulated oocysts after application of various cleaning agents have been examined experimentally by Frenkel and Dubey (5). High concentrations of ammonia (33%) and iodine (7%) killed all oocysts within 30 minutes, whereas ammonia (1.7%) and iodine (20%) were

no more effective than drying alone. Household ammonia (4–6%) requires more than 60 minutes of exposure to sterilize cat feces. Smaller particles of fecal matter (empty, uncleaned litter pan) exposed to water heated to boiling and poured into the pan (75–80°C) are sterilized rapidly.

Despite strong experimental evidence, a direct relationship between oocyst shedding and human toxoplasmosis is not clear from epidemiologic studies. Although 14% to 50% of cats in the United States are seropositive, oocysts are found in the feces of less than 1% of cats examined, perhaps due to the limited age range and short duration (1–3 weeks) of oocyst shedding (19,20). Likewise, the presence of oocysts in the soil correlates poorly with seropositivity in humans (19,20). Additionally, serologic studies of veterinarians and cat owners do not support a strong relationship between a long, intensive exposure to cats and a greater risk of acute toxoplasmosis (21–23). Indeed, persons who report no contact with cats have seropositivity rates of 5% to 40% (21–23).

The latter discrepant information raises questions concerning the role of direct contact with cats and acute toxoplasmosis and our present methods of ascertaining the association. The correlation between seropositivity and house cats may be too crude to reveal a relationship. Studies limited to exposure to sick, young, or feral cats may identify a relationship.

On the other hand, oocysts may be so widely distributed in the environment and direct contact with a cat that is shedding oocysts is so rare that the risk of acute toxoplasmosis through oocysts is a function of age and nonspecific contact, such as outdoor gardening, culinary habits, food preparation, presence of insect vectors, and personal hygiene.

Table 1 Seroprevalences of *Toxoplasma gondii* Among Various Animals in Developed Countries (2)

Animal	Seroprevalence
Cattle	15–50%
Fattening pigs (intensive farming)	1–31%
Pigs	1–21%
Chickens	6–9%
Sheep	16–40%
Goats	19–47%
Horses	1–17%
Dogs	25–40%
Rabbits	6–20%
Pet cats	22–68%
Feral or stray cats	33–68%
Wild turkeys (United States)	17%
Wild horses	16–37%
Deer	30–70%
Moose	15%
Bears (United States)	30–80%
Human (United States) (3)	18–29%

Figure 1 Life cycle of *Toxoplasma gondii*.

The tachyzoite form of the parasite is seen in the acute stage of the infection. Tachyzoites result from ingestion of oocysts, that is, food contamination or tissue cysts (bradyzoites/cystozoites), that is, infected food source. The liberated parasites can survive for 1 to 3 hours in pepsin–hydrochloric acid (stomach fluid) and 4 to 6 hours in trypsin (intestinal fluid)—enough time for transmural migration into the host. The tachyzoites are spread hematogenously and can be identified by Wright's or Giemsa's stains. Every cell type is susceptible to penetration.

The tachyzoite requires an intracellular habitat to survive. The intracellular, intravascular tachyzoite undergoes multiplication, endodyogeny, every 4 to 6 hours to form rosettes. These intracellular collections (tissue cysts) vary in size, but may contain as many as 3000 organisms and are 200 μm in size. Rupture of the host cell results in continuing infection and tissue destruction. As the host develops an immune response to the parasite, parasitemia and systemic infection with tachyzoites decrease. In their "protected" intracellular cysts, the tachyzoites evolve into bradyzoites (cystozoites).

The latent or chronic stage is manifested by tissue cysts in most organs. The brain, skeletal muscle, cardiac muscle, and placenta are the most common sites for tissue cysts, which may be identified by periodic acid-Schiff stain. Recurrent parasitemia is rare in immunocompetent individuals. However, under immunodepressed states, especially AIDS, the cysts may reactivate and cause parasitemia and tissue destruction.

The ingestion of tissue cysts in undercooked or uncooked meat is an important source of infection in the United States. Serologic evidence of past infection and potential for tissue cyst inoculation (2,24,25) is present in 15% of cattle, 15% of sheep, 30% of pigs, and 43% of goats. The reported frequency of tissue cysts in common cuts of meat is less frequent: fresh beef, less than 1%; lamb, 5%; mutton, 10%; and pork, 30% (2,24,25). However, these studies are based on older studies and may underestimate the risk. The selected samples used were too few and too small to estimate the prevalence accurately. It is also probable that modern farming techniques, intensive pen raising, and processed commercial feed reduce the risk. However, the recent movement back to local sources of meat, "reducing your carbon footprint," may lead to more beef, pork, lamb, or chicken that has been exposed to the barnyard cat and its feces.

There is clear evidence to support the relationship between infected meat ingestion and acute toxoplasmosis. Butchers and meat handlers have a higher seropositivity rate than does the general population, through oral ingestion or inoculation of open wounds. In 1965, Desmonts (26) reported a definitive study supporting the meat-to-human hypothesis. Seroconversion rates rose in response to adding undercooked mutton (as a part of therapy) to the diets of children in a French sanitarium. More recently, acute toxoplasmosis was associated with ingestion of undercooked hamburger (27).

Experimental data suggest that attention to food preparation will reduce transmission of tissue cysts. Cleansing food preparation surfaces and utensils with warm soapy water after meat preparation will limit contamination of other foodstuffs. Salting (2% NaCl/liter for 24 hours), drying (24 hours at 18–20°C), freezing (–15°C for 12–24 hours), or heating (to 60°C for 15–20 minutes) sterilizes the meat (28).

Clinical Presentations

Acute Infection

Toxoplasmosis is asymptomatic or very mild (lymphadenopathy and/or fatigue) in 80% of immunocompetent individuals. Cervical, suboccipital, supraclavicular, axillary, or inquired lymph nodes are commonly enlarged. An enlarged single posterior cervical node is highly suspicious. In cases associated with a virulent strain of *T. gondii* or in cases associated with immunosuppressive therapy, AIDS or congenital immunodeficiency syndromes or an immature fetus, clinical signs, or symptoms are present in 30% to 100% of infections.

The clinical syndrome in presumably immunocompetent individuals is best described by an epidemic related to oocyst exposure in a riding stable (29). Of 37 patients with serologically proven toxoplasmosis, 80% had fever, headache, and lymphadenopathy (flu-like syndrome). Myalgia, stiff neck, and anorexia were associated with 50% to 70% of patients. Additionally, arthralgia, rash, mental confusion, and hepatitis were more common than controls. Interestingly, only 3/25 (12%) of patients were correctly diagnosed by their physicians. The combination of lymphadenopathy, fatigue, malaise, atypical lymphocytosis, and/or occasional liver/spleen involvement is similar to mononucleosis or Epstein–Barr infections.

The clinical challenge is to differentiate the nonspecific symptoms of acute toxoplasmosis infection from the symptoms of pregnancy or common viral infections. An interesting study by the U.S. multicentered Toxoplasmosis Study Group (30) performed a detailed interview on 131 mothers of infants with proven congenital toxoplasmosis. Recognizing a serious recall bias, these women reported fever or night sweats (27%), flu-like illness or myalgia (26%), headache (11%), or any "infectious illness" (48%). Importantly, 52% reported no maternal illnesses. Similarly, 35% reported no cat or cat litter exposure and 50% reported no exposure to undercooked meat or poor meat preparation hygiene. In a combined fashion, only 48% had exposure to cat litter, undercooked meat, or toxoplasmosis-like illness during pregnancy. These findings suggest that more than 50% of patients may not be amenable to serologic screening based on risk factors or symptoms.

In immunocompromised patients (AIDS, cancer therapy, transplant patients), reactivation of tissue cysts from prior infection leads to local tissue destruction and, ultimately, systemic parasitemia. As the central nervous system is a common site for tissue cysts, the clinical manifestations include central nervous system abscesses, encephalitis, cognitive impairment, headache, and chorioretinitis. Toxoplasma encephalitis occurs in 30% to 40% of AIDS patients (31).

Most pregnant women are immunocompetent, and clinical expression of the infection ("flu" or lymphadenopathy) is mild and occurs in a minority (5–15%). Serologic evidence of maternal infection varies by maternal age and locale (15,32). The age-related incidence is 20% at less than 25 years, 23% at 26 to 35 years, and 35% at greater than 36 years old. Warm, moist climates have high seropositivity rates (40–60%) and cold, dry climates have much lower rates (10–20%). These observations support the theory that toxoplasmosis is related to the ubiquitous distribution of infectious oocysts and/or food contamination with tissue cysts.

Between 45% and 87% of women will be seronegative and at risk for primary infection during pregnancy. Between 0.5% and 2.3% of pregnant women will be described as being at high risk for fetal infection (probable maternal infection) as indicated by maternal seroconversion. Of these, 5% to 18% demonstrates fetal infection documented by neonatal signs or symptoms and/or positive neonatal *T. gondii*-specific IgM titers.

In a study by Zadik et al. (33), the rate of primary infection with toxoplasma was assessed by seroconversion. In that study, 1621 women had paired serum more than 500 days apart. Primary infection was determined by seroconversion alone. In 160 patients (9.9%), initial titers were positive. Only one seroconversion was detected in 2966 women years. The projected rate for primary infections was 0.23 (95% CI 0.0059–1.3) per 1000 pregnancies. This rate of maternal infection is much lower than predicted by combined serologic methods (i.e., high titer, rising titer, etc.). On the other hand, the neonatal infection rate may be higher as it is a primary infection and true infection is better defined.

While standard in vitro tests have shown little, if any, difference in humoral or cellular immunity between pregnant and nonpregnant women, later infection by certain organisms (i.e., varicella, human papilloma virus) may be reactivated during pregnancy. The combination of latent human immunodeficiency virus (or organ transplantation with immunosuppression), latent toxoplasmosis, and pregnancy often leads to maternal and fetal concerns of reactivation of the toxoplasmosis. In AIDS patients, reactivation usually occurs with CD₄ counts <100 cells/mm³. It appears that pregnant women have no greater risk than that defined by low CD₄ counts. Among 48 pregnant women receiving immunosuppressive therapy after organ transplantation, no cerebral toxoplasmosis was recorded. Among 105 HIV-positive women, only one case of cerebral toxoplasmosis occurred with a CD₄ count of 20 cells/mm² (30). Prophylactic treatment to prevent maternal reactivation and vertical transmission is not indicated for CD₄ counts >100 cells/mm³. At CD₄ counts under 100 cells/mm³, the risk is still small, but prophylaxis may be justified, although benefit is not proven. These observations raise a serious question of whether reactivation of latent toxoplasmosis actually occurs in immunocompetent pregnancies. Seroconversion, rather than high or rising titers, may be the major serologic marker for fetal infection.

Experimental infection of rhesus monkeys showed that transmission of *T. gondii* occurred in four of nine (44%) in second-trimester maternal infections and three of nine (33%) in the third trimester (34). Another four fetuses of mothers infected in the third trimester were probably infected, but the organism could not be isolated. The overall infection rate was 61%. Likewise, pregnant cats were orally inoculated with *T. gondii* (17). Among 8 animals euthanized at 10 or 31 days post-inoculation, 7 of 33 (21%) fetuses from all litters had *T. gondii* isolated. Eight additional pregnant cats were allowed to reach parturition and nurse their kittens. Of 40 kittens from all eight litters, 26 (65%) had *T. gondii* isolated from tissues; 25 of 26 kittens were severely ill by 24 days of life. It is apparent that in susceptible species (cats, rhesus monkeys, and man), the rate of intrauterine fetal infection is 40% to 70%.

Fetal infection is a combination of acute and persistent infection due to relative immune incompetence and reduced passive IgG transfer from the mother. Prior to 30 weeks of gestation, the fetal immune system is of little help; IgM, IgG, and cellular responses are limited and functionally immature (35). For example, in proven cases of in utero infection with toxoplasmosis in the first trimester, funicentesis at 20 to 24 weeks demonstrated four out of nine with toxoplasma-specific IgM (36). Placental transfer of maternal IgG is limited by gestational age and target antigen. Immunoglobulin G transport has been noted by 8 weeks, but remains less than 100mg/dL until 20 weeks. Subsequently, the fetal levels of maternal IgG rise until they are greater than maternal levels.

Acute fetal toxoplasmosis results from initial placental involvement and subsequent systemic fetal parasitemia and infection. The most consistent histologic placental finding is chronic inflammation in the decidua capsularis and focal reaction in the villi. Tissue cysts as well as tachyzoites may be seen. Systemic involvement results in splenomegaly, hepatomegaly, jaundice, anemia, thrombocytopenia, chorioretinitis, and encephalitis. Persistent infection due to limited immune response leads to extramedullary hematopoiesis in the liver and decreased plasma protein production. When coupled with cardiac involvement, nonimmune hydrops may result. Widespread tissue destruction manifests as clinically apparent growth restriction. The most characteristic damage in the central nervous system is extensive necrosis of brain parenchyma secondary to vasculitis; microcephaly is a common sequela. Periventricular and periaqueductal vasculitis is common in toxoplasmosis. Subsequent calcification of these areas is apparent on roentgenographic or ultrasonographic examinations. Obstruction of the aqueduct and/or meningeal involvement leads to hydrocephalus. Chorioretinitis may be present at birth or may develop in the first two decades after birth through recrudescence or persistence of the disease.

Table 2 describes the incidence of organ system involvement in 210 of congenitally infected neonates who were followed prospectively for 1 year (37). Twenty-one (10%) of the infants (including one death) had severe congenital toxoplasmosis

Table 2 Neonatal Toxoplasmosis (n = 210)

Prematurity (≤2500g)	4%
Intrauterine growth restriction	6%
Jaundice	10%
Hepatosplenomegaly	4%
Thrombocytopenia	2%
Anemia	4%
Eosinophilia	2%
Microcephaly	5%
Hydrocephalus	4%
Hypotonia	6%
Seizures	4%
Psychomotor retardation	5%
Intracranial calcifications	11%
Abnormal cerebrospinal fluid	35%
Micro-ophthalmia	3%
Bilateral chorioretinitis	6%
Unilateral chorioretinitis	16%
Normal	55%

Source: From Ref. 37.

with central nervous system involvement. Seventy-one (34%) had a normal clinical exam except for peripheral retinal scarring or isolated asymptomatic intracranial calcifications. One hundred and sixteen (55%) were normal. Caution should be exercised when evaluating the incidence of injury. Most series are populations of referred neonates and represent a population with more clinically apparent diseases. Unless all pregnant women and neonates are screened and all infected neonates are systematically examined, the risk is lower than that published. In addition, women countries, which use universal screening to identify maternal seroconversion and use amniocentesis and PCR to identify fetal infection, are more likely to initiate antibiotic therapy after a significant percentage of women (about 40%) chose to abort (38). Antibiotic therapy may modify (lower) the incidence of severe disease at birth. In the latter study, 58 of 76 (77%) had asymptomatic infection after 84% received antibiotics prior to amniocentesis.

The timing of acute fetal toxoplasmosis greatly affects the sequelae (39) (Table 3). When it occurs early in gestation, especially prior to 13 weeks, the infection is more widespread and destructive. At 11 to 13 weeks, the fetal immune system begins to identify antigens as "self" or "non-self." Toxoplasma antigens that are incorporated into fetal tissue may be misrecognized as "self" and harder to eliminate. In addition, the continuing immaturity of the fetal immune system and limited active transfer of toxoplasma-specific maternal IgG prior to 25 weeks poorly control the infection. Additionally, immune "protected" location, the central nervous system, and inner eye allow the organisms to avoid host responses.

Chronic Infection

Early reports of follow-up indicated that children with clinically apparent disease at birth had a poor prognosis: mental retardation, 93%; seizures, 81%; spasticity or palsies, 70%; impaired vision, 61%; hydrocephalus or microcephaly, 32%; deafness, 15%; and normal, 11% (40). Fortunately, many infants with congenital toxoplasmosis are completely normal at birth (40–55%). The prognosis in mildly affected or asymptomatic infection appears to be much better (Table 2).

A continued risk of damage is apparent in 20-year follow-up of congenitally infected children and in epidemiologic studies. Koppe et al. (41) followed 11 children for 20 years. One of five asymptomatic and treated infants and four of six (two with severe disabilities) symptomatic infants developed new chorioretinitis by 20 years. Wilson et al. (42) followed 23 children with subclinical infections at birth for 1.25 to 17.25 years. Of 23 children, 19 (83%) had one or more episodes of chorioretinitis, 11 (48%) had unilateral or bilateral

blindness, and 12 (53%) had a new diagnosis of central nervous system disorders. Sever et al. (32) performed a multivariate analysis of 22,845 patients in whom toxoplasmosis titers were correlated with poor outcomes during a standardized exam at 7 years of age. The results were controlled for race, maternal age, and socioeconomic index. Visual impairment, microcephaly, coordination problems, right and left identification, and soft neurologic signs correlated with toxoplasmosis titer. Bilateral deafness, extraocular movement, and IQ <70 correlated with the interaction variable, toxoplasmosis titer $\times$ maternal age.

A 20-year prospective study (1985–2005) of 676 pregnancies and 681 fetuses gives more reassuring outcomes than prior data (43). Mothers who were IgG seronegative were followed with monthly serology. Antenatal therapy with spiramycin 9×10^6 SI units per day was initiated at recognition of maternal IgG seroconversion until an amniocentesis (4 weeks later) was performed to identify fetal infections by PCR. After definitive diagnosis of fetal infection, a regimen of sulfadoxine (1000mg/wk), pyrimethamine 950mg/week, and folinic acid 50mg/week was initiated until delivery. The same regimen was used in pediatric doses until 2 years of age. They reported that 17% (112 of 666) live births had congenital toxoplasmosis after maternal seroconversion during pregnancy (monthly IgG serology). Of these, 74% were asymptomatic, 1.5% had severe manifestations at birth, and 26% were diagnosed with chorioretinitis; 6 of 28 infants with chorioretinitis had macular disease (unilateral). Two children had additional neurologic issues, seizures, and psychomotor delay. The partial improvement in outcomes from earlier data suggests a benefit to early recognition for fetal infection and fetal therapy.

Diagnosis

Behavioral Screening

Ideally, all women who are at risk for pregnancy or are trying to get pregnant need assessment and education concerning the risk of toxoplasmosis during pregnancy, that is, preconceptional counseling. Each woman should be asked about her ongoing exposures to potentially contaminated soil or water, cat exposures, kitchen and personal hygiene, and predilection to eating and undercooked meats or game. In high-risk women (any positive response), a preconceptional measurement of toxoplasma-specific IgG will identify potential risk of fetal infection during a subsequent pregnancy. If the toxoplasmosis-specific IgG is positive, then the risk of fetal toxoplasmosis is very small in an immunocompetent individual. If patient is seronegative, a program of primary prevention education is critical (Table 5). During pregnancy, monthly serology will be

Table 3 Time of Maternal Infection and the Frequency of Fetal Infection

	Time of maternal infection		
	First trimester <i>n</i> = 126	Second trimester <i>n</i> = 246	Third trimester <i>n</i> = 128
No infection	86%	71%	41%
Perinatal death	5%	2%	0%
Fetal infection	9%	27%	59%
Severe	6%	2%	0%
Mild	1%	5%	6%
Subclinical	2%	20%	53%

Source: Adapted from Ref. 39 (Table 31.5).

helpful in the early identification of fetal infection and initiation of antibiotics.

Serologic Screening

Maternal antibody response is normal to a toxoplasmosis infection; IgM appears first and rises sharply for 1 to 2 weeks and is sustained for 4 to 6 months (IgM-IFA) or 6 to 12 months (or years) when measured by more sensitive methods [double-sandwiched IgM enzyme-linked immunosorbent assay (IgM-ELISA)]. After the start of the increase in IgM, IgG rises slowly for 1 to 3 weeks (2–50 IU/mL, 1:10–1:200). Between 3 and 6 weeks, IgG antibodies rise sharply to >400 IU/mL or titers >1:1000. In 2% to 4%, the titers are >1:4000 [indirect hemagglutination test (IHA)] and persist at high levels for years. Again, IgG tests vary in their sensitivity.

The majority of infected pregnant women are asymptomatic, and a combination of testing for immunoglobulin G (IgG), M (IgM), and IgA antibodies is used to diagnose toxoplasmosis. These tests include the Sabin-Feldman dye test (dye test), complement fixation, IHA, indirect fluorescence antibody test (IFA), agglutination test, double-sandwiched IgM-ELISA, and the IgM immunosorbent agglutination assay (IgM-ISAGA).

Although the Sabin-Feldman (dye test) is the time-honored gold standard, its technical complexity (mouse inoculation and observation of a biologic response) limits application outside of research settings. In the IHA test, red blood cells tagged with toxoplasma antigen agglutinate when exposed to serum containing IgG and IgM antibodies to toxoplasma. This response occurs several months after an infection. As a result, as many as 50% of infants with fetal infection will have negative IHA in the first 6 months of life. Complement fixing antibodies appear later than dye test antibodies and, depending on the antigen employed, the results vary. The agglutination test uses parasites fixed in formalin and is very sensitive to IgM antibodies. Nonspecific agglutination may occur due to “naturally” occurring IgM (in mothers negative for toxoplasma-specific IgG and IgM antibodies). Unless the laboratory corrects for these antibodies, false positives can occur. The IFA test uses slide preparations of fixed toxoplasma that are washed with the patient’s serum at different dilutions and cross-reacted with fluorescein-tagged antiserum prepared against IgG (anti-IgG). Although the IFA test is equal to the dye test in specificity, patients with connective tissue disorders, that is, positive rheumatoid factor, may test false positive. Positive tests should be confirmed with a dye or IHA test.

The IgM titer is measured in three ways: IgM-indirect fluorescent antibody (IgM-IFA), IgM-ELISA, and IgM-ISAGA. IgM-IFA relies on antiserum specific to IgM. A titer >1:160 is diagnostic of an acute infection. If the antiserum cross-reacts with IgG, which is true of some testing kits, the results may be false positive. Additional problems include an inhibitory effect of high maternal IgG titers on fetal IgM titers and false positives created by autoimmune antibodies, that is, rheumatoid factor.

The conventional IgM-ELISA may also be false positive in patients with chronic infection or autoimmune disease. As a result, Naot et al. (44) developed a double-sandwiched IgM-ELISA. In a direct comparison with IgM-IFA in 55 serum samples from neonates with proven toxoplasmosis, the

double-sandwiched IgM-ELISA was more likely to be positive (81% vs. 25%) than was IgM-IFA.

The IgM-ISAGA combines the advantages of direct agglutination and double-sandwiched ELISA. The IgM-ISAGA does not use an enzyme conjugate and avoids cross-reactivity with rheumatic factor. It is more sensitive than the IgM-IFA or the IgM-ELISA and will be positive 1 to 2 weeks earlier than the latter tests.

Continual improvement in the specificity and sensitivity of toxoplasma-specific IgM testing has muted its usefulness. The more sensitive the test, the less likely it is to narrow the time of the maternal infection if preconceptual Toxoplasmosis IgG testing has not been performed. In the latter situation, the timing is critical. The earlier the infection, especially prior to 14 weeks, the less likely fetal infection has occurred, but it is more likely that severe disease will be manifest at birth. Diagnosis of fetal infection prior to 20 weeks allows the mother a more educated choice whether to carry or abort the pregnancy. In addition, earlier therapy may improve the efficacy of antibiotic therapy.

Commercial testing, the avenue for most clinicians in the United States, can create confusing and difficult decisions (45). The laboratory testing companies are most interested in producing a reliable result with the largest profit margin. Unless the laboratory pre-absorbs the naturally occurring anti-IgG IgM (rheumatoid factor), false positive, low levels of IgM are identified and reported. This occurs more often when the IgG is very high. Overreading the results may create enough concern by the mother to terminate her pregnancy when no fetal infection is present; one study suggests this occurs in 17.5% of women misinformed about the risk of infection (46). Retesting the same serum using different IgM methodologies and identifying other elevated immunoglobulins (IgA, IgE), an abnormal differential agglutination test, or low IgG avidity can identify true positives and reduce the risk of abortion to 0.4% (47). Most commercial laboratories do not save the serum and are reluctant or are inexperienced in the performance of confirmatory testing. The American College of Obstetricians and Gynecologists and the Centers for Disease Control and Prevention recommend contacting a reference laboratory prior to an intervention as a result of the primary serologic screening test. Other useful contact information includes reference laboratory for serology, isolation, and PCR assay (650-853-4828), Congenital Toxoplasmosis Study Group (U.S.), and Educational information on the Internet www.toxoplasmosis.org.

Seroconversion of IgG in a prior seronegative provides unequivocal evidence of infection occurring in the period of time between the two tests. However, most women become pregnant without knowledge of their antibody status and diagnosis must rely on a skilled and experienced interpretation of positive IgG serology, a positive IgM serology, rising antibody titers, or, occasionally, persistent, high titers. Unfortunately, the numerous tests used to measure antibody response have considerable inter- and intra-test variability.

Fetal Diagnosis

The advent of sophisticated ultrasound diagnosis and invasive fetal testing (funicentesis/cordocentesis) has heralded new options in the diagnosis and management of toxoplasmosis. Although serologic testing may identify 0.5% to 2.5% of

pregnant women at high risk of fetal toxoplasmosis, fetal infection is not an absolute diagnosis without measurement of fetal immunologic response or demonstration of the organism. By using advanced ultrasound diagnosis and invasive fetal testing, 5% to 20% of maternal infections in the first 20 weeks of pregnancy and 40% to 80% in the second 20 weeks of pregnancy result in fetal infection. The rapid identification of the fetal infections is the best management.

High-resolution obstetric ultrasound has a role in screening and diagnosis of fetal infection. The fetal findings include ventriculomegaly, microcephaly, intracranial calcifications, increased placental thickness, parenchymal liver calcifications, fetal hydrops, ascites, hyperechoic bowel, and fetal growth restriction. The presence of any one or a combination of these findings in the presence of positive serology has a sensitivity of 49%, specificity of 99%, positive predictive value of 98%, and a negative predictive value of 89% (48). The unanticipated identification of these findings should initiate toxoplasma-specific IgG and IgM serology testing. If positive serology is positive, an amniocentesis for AF-PCR testing is warranted.

Access to fetal tissues (amniotic fluid, amniocytes, chorionic villi, or fetal blood) allows unequivocal fetal diagnosis. In utero fetal infection may be diagnosed through demonstration of toxoplasma antigens (parasites or DNA), fetal immune response (IgM), or physical or laboratory evidence of fetal infection. Inoculation of mice or tissue cultures with fetal specimens is 50% to 70% sensitive in documenting fetal infection. Polymerase chain reaction identification of toxoplasma DNA significantly reduces the time and laboratory requirements for antigen detection. However, adequate laboratory support is essential, but currently there is limited access and availability.

The demonstration of a fetal IgM through cordocentesis is limited by gestational age. Prior to 24 weeks, the upper age limit for abortion in the United States, less than half of fetuses, will mount an IgM response. Abnormal laboratory findings on fetal blood include elevated white cell count, eosinophil count, monocytes, gamma-glutamyl transpeptidase, and lactic dehydrogenase activity (Table 4) (36,49).

Isolation of the parasite from the mother or infant provides unequivocal proof of infection. The parasite (bradyzoite or tachyzoite) can be isolated from blood, muscle, brain,

amniotic fluid, or placental tissue through inoculation of mice or tissue culture. The frequency of isolation from the placenta varies by the trimester in which toxoplasmosis was acquired and untreated: first trimester, 25%; second trimester, 54%; third trimester, 65%; and overall, 50% (50). However, the placenta may be negative in the presence of a congenitally infected fetus. Parasitemia in infected newborns varies by clinical stage: acute/generalized, 71%; neurologic or ocular, 17%; and subclinical, 52% (51). When the diagnosis is attempted by funicentesis in the second trimester, the fetal blood is positive in 70% of cases (52). Overall (term and preterm), the amniotic fluid sediment will contain toxoplasma in 60% of samples from fetuses with congenital infection. Although isolation of the organism is a powerful investigational tool, it requires time and specialized laboratories not readily available to the practicing physician.

Molecular genetics has provided a powerful new tool in antigen identification, the polymerase chain reaction (PCR) (52,53). In 43 cases, PCR was compared with mouse inoculation (tissue/amniotic fluid cultures) and IgM-ISAGA (IgM immunosorbent agglutination) on fetal blood. PCR correctly diagnosed toxoplasmosis as positive in 8 out of 10 samples with no false-positive diagnosis. More recently, Hohlfeld et al. (52) compared a competitive PCR test of amniotic fluid with conventional diagnostic methods (ultrasonography, fetal blood sampling, and serologic diagnosis) in 339 women who had acquired toxoplasmosis during pregnancy (combined serologic diagnosis). Conventional methods identified 34 of 37 (sensitivity 89.7%) neonates with serologically or pathologically confirmed toxoplasmosis at birth. The PCR test identified 36 of 37 infants (sensitivity 97.4%). In a more recent analysis, real-time PCR was 92% sensitive and had a 98% negative predictive value in a study 377 mothers with seroconversion during pregnancy (54). The PCR test on amniotic fluid is now the standard method for the diagnosis of fetal toxoplasmosis.

Universal Screening

The frequency of maternal (0.8%) and fetal (0.32%) infection, the frequency of long-term disability after fetal infection (50–90%), and the cost of caring for those disabilities (\$80,000–\$150,000 per case) describe the potential benefits of a universal screening program. This potential benefit compares favorably

Table 4 Antenatal Diagnosis of Fetal Toxoplasma			
	Daffos et al. (36)	Foulon et al. (49)	Total
Number of high-risk mothers	746	50	796
Fetal infection	42	6	48 (6%)
Diagnosed in utero	39/42	5/6	44/48 (92%)
Positive specific IgM	9/42	1/3	10/45 (22%)
Elevated total IgM	22/37	—	59%
Elevated WBC	16/39	—	41%
Eosinophilia	8/33	2/3	10/36 (28%)
Elevated LDH	7/35	3/3	10/38 (26%)
Elevated GGT	24/34	3/3	27/36 (75%)
Abnormal ultrasound: ascites	2/42	0/3	2/45 (4%)
Ventricular dilation	17/42	1/3	18/45 (40%)
Intracranial calcification	—	1/3	33%
Abbreviations: LDH, lactic dehydrogenase; GGT, gamma-glutamyl transpeptidase.			

with other universal screening programs: syphilis, gonorrhea, phenylketonuria, or maternal serum alpha-fetoprotein. Commercial laboratories would easily and happily accommodate the demand for serologic testing in a universal screening program. Finally, respected researchers endorse the concept of universal screening (55).

On the other hand, the cost, unreliability in testing, and the inadequacy and cost of resources to interpret, diagnose, and treat fetal infection are major obstacles to implementation. The immunofluorescent antibody test, which is used in 80% of commercial laboratories (56), costs \$80 to \$120. Each patient would require two to three tests: two tests to document seroconversion and three tests (in 40% of patients) to document a rising titer (paired samples or IgM). Universal serologic screening would cost an estimated \$320 to \$720 million.

A cost benefit analysis based on data from a Finnish prospective study (57) and on Finnish cost data was performed to compare the no-screening and screening alternative for primary toxoplasma infection during pregnancy (58). Given a 20.3% seropositivity and an incidence of fetal infection to be 2.4/1000 seronegative pregnancies, the total annual costs of congenital toxoplasmosis without screening were \$128 per pregnancy per year (U.S. dollars). With screening, the cost would be \$95. The authors concluded that universal screening, plus health education, is cost effective if the incidence of maternal primary infection exceeds 1.1/1000 pregnancies and the effectiveness of treatment is greater than 22%.

The diagnosis of fetal infection in high-risk mothers is necessary unless the practitioner and patient are willing to abort or treat 9 to 15 uninfected fetuses for each infected fetus. Amniocentesis and/or funicentesis would be needed in 296,000 ($0.8\% \times 3.7$ million) women per year, with each ultrasound-guided procedure costing a minimum of \$1500. The necessary expertise, laboratory support, and financial resources are not available for this demand.

In summary, while selected preconceptional screening for seronegativity and antenatal screening for seroconversion in behaviorally high-risk individuals may be warranted, first-trimester universal screening for toxoplasmosis is not warranted at this time in the United States. However, serologic screening should be considered when the patient has a "flu"-like syndrome, and/or has lymphadenopathy, or when the fetus, on ultrasonographic examination, has growth restriction, nonimmune hydrops, ascites, hydrocephalus, microcephaly, or intracranial calcification.

Antibiotic Therapy

Pyrimethamine, a substituted phenylpyrimidine antimalarial drug, cures toxoplasmosis in laboratory animals through folic acid antagonism. Pyrimethamine produces a reversible, dose-related bone marrow depression. Thrombocytopenia, with its associated bleeding sequelae, is the most common toxic affect, but anemia and leukopenia may occur as well. A semi-weekly complete blood cell and platelet count should be obtained while the patient is being treated.

Sulfadiazine (or sulfapyrazine, sulfamethazine, or sulfamerazine) acts synergistically with pyrimethamine with a combined activity against toxoplasmosis much greater than a merely additive affect. Like pyrimethamine, sulfa drugs disrupt

folic acid metabolism in bacterial species. Renal toxicity (crystalluria and hematuria), toxic dermal reaction (Stevens-Johnson syndrome), and blood dyscrasias are reported with sulfa drug therapies. However, these agents are well tolerated in most individuals. In the neonate, sulfa drugs displace bilirubin from albumin, and kernicterus may occur. The risk of neonatal kernicterus from maternal intake of sulfa drugs is quite small.

The evaluation of antibiotic efficacy depends on the time between the onset of infection and initiation of therapy. In human studies, the timing of fetal toxoplasmosis is very difficult to determine because the infection is usually asymptomatic. Experimental inoculation of infra-human primates, rhesus monkeys, and treatment with anti-toxoplasmosis antibiotics provides powerful insight into the pharmacokinetics of pregnancy and the efficacy of the antibiotics.

Schoondermark-Van de Ven et al. (59) established the dosage regimen for pyrimethamine and sulfadiazine in pregnant rhesus monkeys. The distributions of both drugs followed a one-compartment model. The maximum concentrations in serum were 88.7 µg/nL, 0.22 µg/nL for sulfadiazine and pyrimethamine, respectively. The serum elimination half-lives were much greater for pyrimethamine (44.4 hours) than sulfadiazine (5.2 hours).

Subsequent to the latter experiment, Back et al. used the dosage regimen to treat six rhesus monkeys that were infected with *T. gondii* at 90 days of gestation (second trimester) (60). Four similarly infected pregnant monkeys were used as controls. Infection was documented by isolation of *T. gondii* from amniotic fluid samples. The treated animals received pyrimethazine 1 mg/kg body weight and sulfadiazine 50 mg/kg/day for the remainder of the pregnancy. *T. gondii* was present at birth in three of four untreated neonates. In all animals treated with pyrimethazine and sulfadiazine, no parasite was identified in the amniotic fluid 10 to 13 days after initiation of therapy. No neonate demonstrated *T. gondii* at birth.

Spiramycin, a macrolide antibiotic, has an antibacterial spectrum similar to erythromycin and is active against toxoplasmosis in laboratory animals. With spiramycin 2 g/day, the concentrations were maternal serum, 1.19 µg/nL; umbilical cord, 0.63 µg/nL; and placenta, 2.75 µg/nL. With spiramycin 3 g/day, the concentrations were maternal serum, 1.69 µg/nL; umbilical cord serum, 0.78 µg/nL; and placenta, 6.2 µg/nL (59,60). Persistent, high tissue levels account for its greater activity against toxoplasmosis than erythromycin. During pregnancy, spiramycin is concentrated in placental tissue two to four times the serum levels, an important advantage, as placental infection is sentinel to fetal infection. Spiramycin seems to be well tolerated. When used as an antineoplastic agent in doses greater than 35 mg/kg, some patients experienced vertigo, nausea, vomiting, and anorexia. No impairment of the heart, blood, liver, or kidneys was noted (60).

Schoondermark-Van de Ven et al. (61) inoculated eight rhesus monkeys at 90 days of gestation (second trimester). The parasite was found in five of eight amniotic fluids at subsequent sampling. These monkeys with documented fetal infection were treated with intravenous spiramycin 20 mg/kg/day in two divided doses. At 14 days after initiation of spiramycin, *T. gondii* was still detectable by PCR analysis,

but not by mouse inoculation in four of five amniotic fluids. In four monkeys that had received spiramycin for 7 weeks, no *T. gondii* was detected by PCR or mouse inoculation in the placenta, amniotic fluid, or neonatal organs. One fetus that delivered prematurely after 2 weeks of spiramycin had *T. gondii* associated with its placenta based on PCR results alone. Spiramycin accumulated in extra-central nervous system tissues at concentrations 5 to 28 times umbilical cord concentrations; but no spiramycin was found in fetal/neonatal brains. These data suggest that in primates early treatment with spiramycin may eliminate fetal infection with moderate success, but if there is fetal central nervous system involvement, spiramycin is likely to be ineffective.

Clinicians have come to expect that antibiotics will effectively treat infection. Unfortunately, effective antibiotic treatment of fetal toxoplasmosis has not been realized. In the last 10 years, three separate meta-analyses have concluded little benefit from antibiotic therapy. No trials with randomized assignment of women with confirmed fetal toxoplasmosis have been preformed; all are observational studies (62–64). The Systematic Review on Congenital Toxoplasmosis study group systematically reviewed 26 cohorts where spiramycin and/or pyrimethamine–sulfonamide were used to treat fetal toxoplasmosis (64). Overall, treatment did not change the frequency of any clinical manifestation: adjusted odds ratio = 1.11 ($p = 0.74$). In a sub-analysis, there was a small reduction in maternal to fetal transmission when spiramycin was initiated less than 3 weeks after maternal seroconversion: adjusted odds ratio (95th CI) = 0.48(0.28–0.80). Maternal to fetal transmission was strongly associated with gestational age at seroconversion. The odds of transmission increased by 12% per week of maternal gestation; the estimated rate of maternal to fetal transmission was 15% (95th CI 13–17%) at 13 weeks, 44% (40–47%) at 26 weeks, and 71% (66–76%) at 36 weeks.

The findings of early treatment after seroconversion and the results of the earlier quoted study of Berribi et al. (43), provide some guidance for therapy. In summary, a potential algorithm is preconceptual screening (or first prenatal visit) and monthly screening with *Toxoplasma* serology during pregnancy in seronegative women. If seroconversion occurs, then spiramycin initiated until an amniocentesis can be performed for diagnosis of fetal infection. If fetal infection is diagnosed, then treatment with pyrimethamine/sulfonamide and folinic acid is initiated antenatally and neonatally.

Prevention

Primary prevention has the potential to greatly reduce the medical and economic costs of congenital toxoplasmosis. A combination of public education, that is, high school curriculum, preconceptual counseling, and early prenatal counseling, would be most effective in reducing congenital toxoplasmosis. In addition to material concerning the risks of toxoplasmosis, the role of cats and the general guidelines are listed in Table 5.

Foulon et al. (65) evaluated the reduction in the incidence of congenital toxoplasmosis using a primary prevention scheme (education) and a secondary prevention scheme (screening and treatment). Among the 11,286 consecutive pregnant women, primary prevention reduced the seroconversion rate by 63% ($p = 0.013$). Of 76 women determined to be at risk for

Table 5 Primary Prevention of Toxoplasmosis During Pregnancy
<i>Oocyst exposure</i>
Avoid the purchase of a new cat just prior to or during pregnancy
Avoid sick or feral cats
Use gloves for gardening
Have other family members clean the litter box
Clean the litter box daily
Sterilize the empty pan with boiling water for 20 minutes
Prevent flies and other vectors from contaminating food
Thoroughly wash fruits and vegetables
<i>Tissue cyst exposure</i>
Cook meat to 150°F (66°C)
Cleanse the preparation surface immediately after meat preparation
Wash hands immediately after meat preparation
Avoid contact with mucous membranes of the mouth and eyes while handling raw meat

delivering a child with congenital toxoplasmosis, 8 (11%) infected fetuses were detected by prenatal fetal diagnosis. Of these, secondary prevention (antibiotics) reduced the incidence of congenital toxoplasmosis by an additional 40%. Unfortunately, a recent meta-analysis of randomized trials showed poor design quality and no benefit in reducing the incidence congenital toxoplasmosis. Regardless, given the benefit of the Foulon et al. study and the low cost of behavioral interventions, the author recommends their continued use (66).

SUMMARY

The medical costs of maternal toxoplasmosis warrant the development of effective programs in primary prevention, diagnosis, and therapy. Maternal toxoplasmosis occurs in 0.5% to 1% of pregnancies, and 10% to 60% of their fetuses will be infected. While the majority (80–90%) of infected neonates will be asymptomatic at birth, a large proportion (20–90%) will develop one or more disabilities. These include hydrocephalus, microcephaly, chorioretinitis, mental retardation, and deafness.

Primary prevention through education and monitoring of food sources has an unfulfilled potential. The effectiveness of primary prevention is measured by changes in behavior and seroconversion rates. A universal screening program, especially preconceptual screening, will improve diagnosis and allow abortion as an option. However, the lack of standardized tests and poor correlation between maternal infection and fetal infection may result in many unnecessary abortions.

Fetal diagnosis allows more focused therapy. It will significantly reduce unnecessary abortion and antibiotic therapy. Unfortunately, fetal diagnosis requires sophisticated clinical skills, equipment, and laboratory support that are currently available only at major medical centers.

The inability to pinpoint the onset or severity of fetal toxoplasmosis limits the effectiveness of antibiotic regimens used to treat fetal toxoplasmosis. Despite therapy, 10% to 50% of fetuses infected in the first half of pregnancy will manifest injury. This failure rate may be related to an ineffective antibiotic regimen or inadvertently delayed treatment.

Comparative trials of antibiotics are critically important to the management of fetal toxoplasmosis in the future.

REFERENCES

- Luft BJ, Remington JS. Toxoplasmic encephalitis. *J Infect Dis* 1988; 157: 1.
- Tenter AM, Heckworth AR, Weiss LM. *Toxoplasma gondii*: from animal to human. *Int J Parasitol* 2000; 30: 1217.
- Jones JL, Krueszon-Moran D, Wilson M, et al. Toxoplasmosis gondii infection in the United States: seroprevalence and risk factors. *Am J Epidemiol* 2001; 154: 357–65.
- Dubey JP, Miller NL, Frenkel JK. The *Toxoplasma gondii* oocyst from cat feces. *J Exp Med* 1970; 132: 636.
- Frenkel JK, Dubey JP. Toxoplasmosis and its prevention in cats and man. *J Infect Dis* 1972; 126: 664.
- Dubey JP, Hoover EA, Walls KW. Effect of age and sex on the acquisition of immunity to toxoplasmosis in cats. *J Protozool* 1977; 24: 184.
- Dubey JP, Frenkel JK. Experimental toxoplasma infection in mice with strains producing oocysts. *J Parasitol* 1973; 59: 505.
- Dabritz HA, Atwill ER, Gardner IA, Miller MA, Conrad PA. Outdoor fecal deposition by the roaming cats and the attitudes of cat owners and nonowners toward stray pets, wildlife, and water pollution. *J Am Vet Med Assoc* 2006; 229: 74.
- Shapiro K, Conrad PA, Mazet JA, et al. Effect of estuarine wetland degradation on transport of *Toxoplasma gondii* surrogates from land to sea. *Appl Environ Microbiol* 2010; 76: 6821–8.
- Meerburg BG, Kijlstra A. Changing climate-changing pathogens: *Toxoplasma gondii* in North-Western Europe. *Parasitol Res* 2009; 105: 17–24.
- Lass A, Pietkiewicz H, Modzelewska E, et al. Detection of *Toxoplasma gondii* oocysts in environmental samples using molecular methods. *Eur J Clin Microbiol Infect Dis* 2009; 28: 599–605.
- Kappareud G, Jenum PA, Stray-Peterson B, et al. Risk factors for *Toxoplasma gondii* infection in pregnancy. *Am J Epidemiol* 1996; 144: 405–12.
- Cook AJC, Gilbert RE, Buffalano W, et al. Sources of toxoplasma infection in pregnant women: European multicentre case-control study. *Br Med J* 2000; 321: 142–7.
- Boyer KM, Holfels E, Roizen N, et al. Risk factors for *Toxoplasma gondii* infection in mothers of infants with congenital toxoplasmosis: implications for prenatal management and screening. *Am J Obstet Gynecol* 2005; 192: 564–71.
- Feldman HA. Epidemiology of toxoplasma infections. *Epidemiol Rev* 1982; 4: 204.
- Remington JS, McLeod R, Thulliez P, Desmonts G. Toxoplasmosis. In: Remington JS, Klein JO, Wilson CB, Baker CT, eds. *Infectious Disease of the Fetus and Newborn Infant*, 6th edn. Chapter 31. Philadelphia, Pa: Elsevier 2006; Table 31–16: 968–969.
- Dubey JP, Lappin MR, Thulliez P. Diagnosis of induced toxoplasmosis in neonatal cats. *J Am Vet Med Assoc* 1995; 207: 179–85.
- Yilmaz SM, Hopkins SH. Effects of different conditions on duration of infectivity of *Toxoplasma gondii* oocysts. *J Parasitol* 1972; 58: 938.
- Dubey JP. Toxoplasmosis in cats. *Feline Pract* 1986; 16: 12.
- Dubey JP. Toxoplasmosis. *J Am Vet Med Assoc* 1986; 189: 166.
- Behymer RD, Harlow DR, Behymer DE, Franti CE. Serologic diagnosis of toxoplasmosis and prevalence of *Toxoplasma gondii* antibodies in selected feline, canine, and human populations. *J Am Vet Med Assoc* 1973; 162: 959.
- Sengbusch HG, Sengbusch LA. Toxoplasma antibody prevalence in veterinary personnel and a selected population not exposed to cats. *Am J Epidemiol* 1976; 103: 595.
- Ganley JP, Comstock GW. Association of cats and toxoplasmosis. *Am J Epidemiol* 1980; 111: 238.
- Jacobs L, Remington JS, Melton ML. A survey of meat samples from swine, cattle, and sheep for the presence of encysted toxoplasma. *J Parasitol* 1960; 46: 23.
- Krogstad DJ, Juranek DD, Walls KW. Toxoplasmosis: with comments on risk of infection from cats. *Ann Intern Med* 1972; 77: 773.
- Desmonts G, Couvreur J, Alison F, et al. Etude épidémiologique sur la toxoplasmose: de l'influence de la cuisson des viandes de boucherie sur la fréquence de l'infection humaine. *Rev Fr Etud Clin Biol* 1965; 10: 952.
- Kean BH, Kimball AC, Christianson WN. An epidemic of acute toxoplasmosis. *JAMA* 1969; 208: 1002.
- Jacobs L, Remington JS, Melton ML. The resistance of the encysted form of *Toxoplasma gondii*. *J Parasitol* 1957; 46: 11.
- Teutsch SM, Juranek DD, Sulzer A, Dubey JP, Sikes RK. Epidemic toxoplasmosis associated with infected cats. *N Engl J Med* 1979; 300: 695.
- Boyer KM, Holfels E, Roizen N, et al. Risk factors for *Toxoplasmosis gondii* infection of mothers of infant with congenital toxoplasmosis: Implications for prenatal management and screening. *Am J Obstet Gynecol* 2005; 192: 564–71.
- Biedermann K, Flepp M, Fierz W, Joller-Jemelka H, Kleihues P. Pregnancy, immunosuppression and reactivation of latent toxoplasmosis. *J Perinatal Med* 1995; 23: 191–203.
- Sever JL, Ellenbery JH, Ley AC, et al. Toxoplasmosis: maternal and pediatric findings in 23,000 pregnancies. *Pediatrics* 1988; 82: 181.
- Zadik PM, Kudesia G, Siddons AD. Low incidence of primary infection with toxoplasma among women in Sheffield: a seroconversion study. *Br J Obstet Gynecol* 1995; 102: 608–10.
- Schoondermark-Van de Ven E, Melchers W, Galama J, et al. Congenital toxoplasmosis: an experimental study in rhesus monkeys for transmission and prenatal diagnosis. *Exp Parasitol* 1993; 77: 200–11.
- Wilson CB. Developmental immunology and the role of last defense in neonatal susceptibility. In: Remington JS, Klein JO, eds. *Infectious Disease of the Fetus and Newborn Infant*. 3rd edn. Philadelphia PA, W B Saunders, 1990: 17–67.
- Daffos F, Forestier F, Capella-Pavlovsky M, et al. Prenatal management of 746 pregnancies at risk for congenital toxoplasmosis. *N Engl J Med* 1988; 318: 271.
- Couvreur J, Desmonts G, Tournier G, et al. Etude d'une série homogène de 210 cas de toxoplasmose congénitale chez des nourrissons >6 mois et de jupistes de faible hon prospective. *Ann Pediatr* 1984; 31: 815.
- Romand S, Chosson M, Franck J, et al. Usefulness of quantitative polymerase reaction in amniotic fluid as an early marker for fetal infection with *Toxoplasmosis gondii*. *Am J Obstet Gynecol* 2004; 190: 797–802.
- Remington JS, Desmonts G. Toxoplasmosis. In: Remington JS, Klein JO, eds. *Infectious Disease in the Fetus and Newborn Infant*, 3rd edn. Philadelphia, PA: WB Saunders, 1990: 100.
- Eichenwald H. A study of congenital toxoplasmosis. In: Siim JC, ed. *Human toxoplasmosis*. Copenhagen: Munksgaard, 1960: 41–9.
- Koppe JG, Loewer-Sieger DH, deRoever-Bonnet H. Results of 20-year follow-up of congenital toxoplasmosis. *Lancet* 1986; 1: 254.
- Wilson CB, Remington JS, Stagno S, et al. Development of adverse sequelae in children born with subclinical congenital toxoplasma infection. *Pediatrics* 1980; 66: 767.
- Berrebi A, Assouline C, Bessieres MH, et al. Long-term outcome of children with congenital toxoplasmosis. *Am J Obstet Gynecol* 2010; 203: 552.e1–6.
- Naot Y, Desmonts G, Remington JS. IgM enzyme-linked immunosorbent assay test for the diagnosis of congenital toxoplasma infection. *J Pediatr* 1981; 98: 32.
- Newton ER. Diagnosis of perinatal TORCH infections. *Clin Obstet Gynecol* 1999; 42: 59–70.
- Liesenfeld O, Montoya JG, Kinney S, Press C, Remington JC. Effect of testing for IgG avidity in the diagnosis of *Toxoplasmosis gondii* infection in pregnant women: experience in a US reference laboratory. *J Inf Dis* 2001; 183: 1248–53.
- Liesenfeld O, Montoya JG, Tathineni NJ, et al. Confirmatory serologic testing for acute toxoplasmosis and the rate of induced abortions among women reported to have positive *Toxoplasma immunoglobulin M* antibody titers. *Am J Obstet Gynecol* 2001; 184: 140–5.
- Crino JP. Ultrasound and fetal diagnosis of perinatal infection. *Clin Obstet Gynecol* 1999; 42: 71–80.
- Foulon W, Naessens A, Mahler T, et al. Prenatal diagnosis of congenital toxoplasmosis. *Obstet Gynecol* 1990; 76: 769.
- Couvreur J, Desmonts G, Thulliez PH. Prophylaxis of congenital toxoplasmosis: effects of spiramycin on placental infection. *J Antimicrob Chemother* 1988; 22: 193.
- Desmonts G, Couvreur J. Toxoplasmosis: epidemiologic and serologic aspects of perinatal infection. In: Krugman S, Gershon AA, eds. *Infections*

- of the fetus and the newborn infant. Progress in clinical and biological research. Volume 3. New York, NY: Alan R. Liss, Inc., 1975: 115–32.
52. Hohlfeld P, Daffos F, Costa JM, et al. Prenatal diagnosis of congenital toxoplasmosis with a polymerase-chain-reaction test on amniotic fluid. *NEJM* 1994; 331: 695–9.
53. Grover CM, Thulliez P, Remington JS, Boothroyd JC. Rapid prenatal diagnosis of congenital toxoplasma infection by using polymerase chain reaction and amniotic fluid. *J Clin Microbiol* 1990; 28: 2297.
54. Wallon M, Franck J, Thulliez P, et al. Accuracy of real-time polymerase chain reaction for *Toxoplasma gondii* in amniotic fluid. *Obstet Gynecol* 2010; 115: 727–33.
55. Wilson CB, Remington JS. What can be done to prevent congenital toxoplasmosis? *Am J Obstet Gynecol* 1980; 138: 357.
56. Sever JL. TORCH tests and what they mean. *Am J Obstet Gynecol* 1985; 152: 495.
57. Lappalainen M, Koskela P, Hedman K, et al. Incidence of primary toxoplasma infections during pregnancy in southern Finland: a prospective cohort study. *Scand J Infect Dis* 1992; 24: 97–104.
58. Lappalainen M, Sintonen H, Koskiniemi M, et al. Cost-benefit analysis of screening for toxoplasmosis during pregnancy. *Scand J Infect Dis* 1995; 27: 265–72.
59. Schoondermark-Van de Ven E, Melchers W, Camps W, et al. Effectiveness of spiramycin for treatment of congenital *Toxoplasma gondii* infection in rhesus monkeys. *Antimicrob Agents Chemother* 1994; 38: 1930–6.
60. Back N, Ambrus JL, Velasco H, et al. Clinical and experimental pharmacology of parenteral spiramycin. *Clin Pharmacol Ther* 1962; 3: 305.
61. Schoondermark-Van de Ven E, Galama J, Camps W, et al. Pharmacokinetics of spiramycin in the rhesus monkey: transplacental passage and distribution in tissue in the fetus. *Antimicrob Agents Chemother* 1994; 38: 1922–9.
62. Peyron F, Wallon M, Liou C, Garner P. Treatments for toxoplasmosis in pregnancy. *Cochrane Database Syst Rev* 2010: CD001684.
63. Peyron F, Wallon M, Liou C, Garner P. Treatments for toxoplasmosis in pregnancy. *Cochrane Database Syst Rev* 2001: CD001684.
64. Thiebaut R, Leproust S, Chene G, Gilbert R. Effectiveness of prenatal treatment for congenital toxoplasmosis: a meta-analysis of individual patients' data. SYROCOT (Systematic Review on Congenital Toxoplasmosis) study group. *Lancet* 2007; 369: 115–22.
65. Foulon W, Naessens A, Derde MP. Evaluation of the possibilities for preventing congenital toxoplasmosis. *Am J Perinatol* 1994; 11: 57–62.
66. Di Mario S, Basevi V, Gagliotti C, et al. Prenatal education for congenital toxoplasmosis (review). *Cochrane Database Syst Rev* 2009: CD006171.